



Perspectives on setups and experiments
for nuclear data with the gamma-ray
beams at ELI-NP

Pär-Anders Söderström
par.anders@eli-np.ro

P(ND)²-3 Nuclear Data for the Next Decade



- ▶ ELI Beamlines, Prague
- ▶ ELI ALPS, Szeged
- ▶ ELI Nuclear Physics, Măgurele
- ▶ ELI-NP is based on two main components
- ▶ High-power lasers (HPLS) 100 TW, 1 PW, 10 PW, Thales Romania
- ▶ High-brilliance gamma-beam (ELI-GBS)




- ▶ ELI-NP is based on two main components
- ▶ High-power lasers (HPLS) 100 TW, 1 PW, 10 PW, Thales Romania
- ▶ High-brilliance gamma-beam (ELI-GBS)

https://www.eli-np.ro/gsd_second.php

Parameter	Specification
Maximum Photon Energy	19.5 MeV
Tunability of the Photon Energy	Steplessly variable
Linear Polarization of Gamma-Ray Beam	> 95 %
Average Relative Bandwidth of Gamma-Ray Beam (FWHM)	$\leq 5.0 \times 10^{-3}$
Total Photon Flux	$\geq 1.0 \times 10^{11}$ 1/s
Time-Average Spectral Density at Peak Energy	$\geq 5.0 \times 10^3$ 1/(s eV)

Gamma Production System – contract with RI for:


transport line + interaction chamber & optical cavity – finalized by 11/26



	Conservative	Technical limit
Flux (photons/sec)	2.4×10^5	4.3×10^7

C. Matei: private communication


Components for the Upgrade for existing ELI-NP LINAC – contract with RI for: multi-bunch electron source (50 Hz x 200 micro pulses) + x2 accelerator structures, more modulators + klystrons – to be delivered by 11/2026 (installation 2027+)



	Conservative	Technical limit	Development potential
Flux (photons/sec)	2.5×10^7	1.8×10^9	4×10^{11}


C. Matei: private communication

- ▶ Nuclear Resonance Fluorescence spectroscopy using (γ, γ') reactions and the HPGe array ELIADE
- ▶ Pygmy- and giant resonances, level population following neutron emission, and neutron resonances, using (γ, n) reactions in ELIGANT
- ▶ Photo-induced fission cross-sections, fragment distributions, and fission isomers with the ELI-BIC and ELITHGEM instruments
- ▶ Production of exotic isotopes using through a γ -induced ISOL facility
- ▶ Properties of key astrophysics (γ, p) and (γ, α) reactions using ELISSA Si array and ELI-eTPC time-projection chamber
- ▶ High-intensity positron beamline for material research and characterization
- ▶ Applications of Nuclear Resonance Fluorescence for non-proliferation, waste-management and cultural heritage with ELIADE




- ▶ An array of CeBr and LaBr for γ -rays, liquid scintillators and Li-glass detectors for neutrons
- ▶ Tested in-beam (2022-2025 campaigns at ROSPHERE, IFIN 9MV)

P.-A. Söderström, et al.: Nucl. Instrum. Methods Phys. Res. A 1027 (2022) 166171




- ▶ ^3He tube array contained in a polyethylene moderator for neutron counting
- ▶ Detector is operational
- ▶ Tested in-beam




P.-A. Söderström, et al.: Nucl. Instrum. Meth. A 1084 (2026) 171229





P.-A. Söderström, et al.:
Nucl. Instrum. Meth. A
1084 (2026) 171229



- ▶ Thermal neutron capture on ^3He gas
 - ▶ Low γ -ray sensitivity
 - ▶ Integrate total counts
-
- ▶ $28\ ^3\text{He}$ neutron counters
 - ▶ 12 bar gas pressure
 - ▶ Arranged in three rings for neutron energy sensitivity via ring ratio method
 - ▶ Flat efficiency for minimizing systematic uncertainties





- ▶ First campaign of (γ, xn) experiments performed at NewSUBARU LCS beamline, D. M. Filipescu and I. Gheorghe
- ▶ Part of the *Coordinated Research Project on Photonuclear Data and Photon Strength Functions* (IAEA CRP F41032)
- ▶ Wealth of experimental data, continued measurements possible at ELI-NP
- ▶ *T. Kawano, et al., IAEA Photonuclear Data Library 2019, Nucl. Data Sheets, 163 (2020) 109*
- ▶ *S. Goriely, et al., Reference database for photon strength functions, Eur. Phys. J. A55 (2019) 172*

D. M. Filipescu, et al., Phys. Rev. C 90 (2014) 064616

I. Gheorghe, et al., Phys. Rev. C 96 (2017) 044604




- ▶ ELI-NP; Nuclear Physics Department; Applied Nuclear Physics Department at IFIN-HH
- ▶ National Physical Laboratory (NPL), UK
- ▶ Indian institute of Technology Ropar, India
- ▶ UK Atomic Energy Authority (UKAEA), UK

- ▶ Complementary program to measure (α, n) cross-sections in the energy range of the α decay from actinides
- ▶ Mainly energy range of interest 3 – 6 MeV, but other ranges for specific applications
- ▶ 3 MV and 9 MV Tandatron accelerators at IFIN-HH
- ▶ Example 1: ^{19}F (α, n) induced neutron background from ^{234}U in UF_6
- ▶ Example 2: ^{27}Al (α, n) induced neutron background in low-background experimental constructions


R. Roy: Ph.D. Thesis, Indian Institute of Technology Ropar (2025)

$^{27}\text{Al}(\alpha, n)$




R. Roy, et al.: *Phys. Rev. C*, 112 (2025) 044613

$^{19}\text{F}(\alpha, n)$




R. Roy, et al.: *in manuscript*

Neutron energy spectrum




Example of the impact of the $^{19}\text{F}(\alpha, n)$ cross-sections using three different data sets


- ▶ The cross-sections were included in the Geant4 based (α, n) source simulation code SaG4n
E. Mendoza, et al.: *Nucl. Instrum. Meth. Phys. Res. A* 960, 163659 (2020)
- ▶ ^{234}U α source isotropically distributed in a UF_6 cylinder of 76 cm diameter and 206 cm length




- ▶ Next step: Upgraded version ELIGANT-TN-74
- ▶ Add 44 new detectors
- ▶ Four-ring concept for improved ring ratios
- ▶ Extend the flat-efficiency region
- ▶ Collaboration between ELI-NP/IFIN-HH and NCSR Demokritos



H.T. Aslani: private communication




H.T. Aslani: private communication



- ▶ An array of CeBr and LaBr for γ -rays, liquid scintillators and Li-glass detectors for neutrons
- ▶ Tested in-beam (2022-2025 campaigns at ROSPHERE, IFIN 9MV)

P.-A. Söderström, et al.: Nucl. Instrum. Methods Phys. Res. A 1027 (2022) 166171




- ▶ ^3He tube array contained in a polyethylene moderator for neutron counting
- ▶ Detector is operational
- ▶ Tested in-beam


P.-A. Söderström, et al.: Nucl. Instrum. Meth. A 1084 (2026) 171229

- ▶ Excitation cross section measured with polarized protons at RCNP – Good case for commissioning
- ▶ No information on the GDR decay to ground or high lying states through gamma or neutron emission – Good case for day-1
- ▶ Clean measurement of the absolute value of GDR ground state γ -decay
- ▶ Measure the energy dependence of the GDR $B(E1)$
- ▶ Simultaneous $\sigma(\gamma, \gamma)$ and $\sigma(\gamma, n)$ to extract details of the wave function
- ▶ Finer structure from higher-order (3p-3h,...) coupling?
- ▶ What is the nature of the PDR?

F. Camera, et al., *Rom. Rep. Phys.* 68 (2016) S539 M. Krzysiek, et al., *Nucl. Instrum. Meth. A* 916 (2019) 257 P.-A. Söderström, et al., *Nucl. Instrum. Meth. A* 1027 (2022) 166171




A. Tamii, et al.: *Phys. Rev. C* 107, 062502 (2011)



The concept of nuclear photon strength functions:
A model-independent approach via ($\vec{\gamma}$, $\gamma'\gamma''$) reactions

J. Isaak^{a,b,c,*}, D. Savran^b, B. Löher^{a,b}, T. Beck^a, M. Bhike^d, U. Gayer^a, Krishichayan^d,
N. Pietralla^a, M. Scheck^e, W. Tornow^d, V. Werner^a, A. Zilges^f, M. Zweidinger^a


J. Isaak, et al.: Phys. Lett. B 788, 225 (2019)





- ▶ γ -ray strength functions can be measured in a model independent way with γ beams
- ▶ Ratio method ($k, j \neq 0$)

$$\frac{\sigma_{ik}}{\sigma_{ij}} = \frac{f(\Delta E_{ik})}{f(\Delta E_{ij})} \frac{\Delta E_{ij}^3}{\Delta E_{ik}^3} \quad (1)$$




- ▶ High-resolution beams can clearly separate different states!








- ▶ 34 $\text{LaBr}_3\text{:Ce}$ and CeBr_3 detectors
- ▶ 36 EJ-301 liquid scintillator detectors
- ▶ 25 GS-20 lithium glass detectors

P.-A. Söderström, et al.: Nucl. Instrum. Meth. A 1027 (2022) 166171








- ▶ ELIGANT instrumentation to investigate “legacy sources”
- ▶ Subtracting tagged from untagged gives direct n transition from excited states in ^{13}C to the ^{12}C ground state
- ▶ Peak like structures originate from known resonances in ^{13}C
- ▶ Compare the results with source simulation software using latest cross-section libraries
E. Mendoza, et al.: Nucl. Instrum. Methods Phys. Res. A. 960, 163659 (2020)
- ▶ Underestimate the low-energy resonances and overestimate high-energy branching to first excited state in ^{12}C





P.-A. Söderström, et al.: Appl. Radiat. Isot., 167 (2021) 109441



- ▶ Simple angular correlations works well
- ▶ Low-energy (250 keV) - high-energy (>1.25 MeV) neutron angular correlations from Li-glass detectors
- ▶ Can obtain the three-neutron angular correlations
- ▶ Assume neutrons emitted in a plane \perp to the spin axis J :
- ▶ Angular distribution of third neutron relative to J




A. Chietera, et al.:
Eur. Phys. J. A 54, 98
(2018)




D. Choudhury, et al.: *in manuscript*
Three-neutron angular correlations;
 $\theta_{2,3}$ given $\theta_{1,2}$

- Experimental setup to measure the prompt fission neutrons and γ -rays in coincidence with fission fragment



ELIGANT array at ELI-NP facility




Inside view of the vacuum chamber

Data taking duration : 6 months


Holder for ^{252}Cf source mounting

Double sided silicon strip detector (DSSSD) for fission fragment detection


S. Dhuri, et al.: private communication




S. Dhuri, et al.: private communication






- ▶ Combining the large volume γ -ray detectors with the ROSPHERE anti-Compton shields
- ▶ In-beam experiments using the 9MV Tandem at IFIN-HH
- ▶ Collaboration between ELI-NP and Department of Nuclear Physics
- ▶ Clean measurements of high-energy γ -rays



S. Aogaki, et al.: Nucl. Instrum. Methods Phys. Res. A 1056 (2023) 168628






K. Sakanashi, et al.: Phys. Lett. B, 870 (2025) 139893, S-R. Ban, et al.: UPB Sci. Bull. A, 87(3) (2025) 165







- ▶ Going towards fully digital, self-triggering, DAQ
- ▶ Increase counting rate a factor of 50 for Tandem experiments
- ▶ Implement complex pulse-shape discrimination procedures directly on the FPGA

S. Aogaki, et al.: IEEE Trans. Nucl. Sci., 72 (2025) 3652



A. Kuşoğlu.: Private communication

First experiment: results from the Sn nuclei




P.-A. Söderström, et al.: *Phys. Rev. C* 112 (2025) 024327, Calculations: N. Tsoneva

The concept of nuclear photon strength functions:
A model-independent approach via ($\bar{\gamma}$, $\gamma'\gamma''$) reactions

J. Isaak^{a,b,c,*}, D. Savran^b, B. Löher^{a,b}, T. Beck^a, M. Bhike^d, U. Gayer^a, Krishichayan^d,
N. Pietralla^a, M. Scheck^e, W. Tornow^d, V. Werner^a, A. Zilges^f, M. Zweidinger^a


J. Isaak, et al.: Phys. Lett. B 788, 225 (2019)







- ▶ γ -ray strength functions can be measured in a model independent way with γ beams
- ▶ Ratio method ($k, j \neq 0$)

$$\frac{\sigma_{ik}}{\sigma_{ij}} = \frac{f(\Delta E_{ik})}{f(\Delta E_{ij})} \frac{\Delta E_{ij}^3}{\Delta E_{ik}^3} \quad (2)$$

- ▶ High-resolution beams can clearly separate different states!








- ▶ Normalize γ SF on the (γ, γ') data from HI γ S
- ▶ Use the normalization to fix the NLD slope

P.-A. Söderström, et al.: Phys. Scr. 100, 075301 (2025)



- ▶ All normalisation methods give consistent results
- ▶ No (or very small) violation of the Brink-Axel hypothesis observed!
- ▶ Several different complementary experimental methods and approaches available in the same campus!

P.-A. Söderström, et al.: In manuscript


Nuclear data for Ultra High-Energy Cosmic Rays



Nuclear data for Ultra High-Energy Cosmic Rays



T. Inakura, et al.: Phys. Rev. C 84, 021302(R) (2011)



M. Kimura, Y. Taniguchi:
Private communication

Astroparticle physics



Japan



Romania



South Africa

Theory

- ▶ Propagation Calculations
- ▶ Galactic magnetic field
- ▶ Acceleration mechanisms

Experiment

- ▶ Pierre Auger
- ▶ Telescope Array
- ▶ Source direction
- ▶ Composition

Theory

- ▶ Reaction theory
- ▶ Structure theory
- ▶ Shell model, mean-field models, ab-initio models...

Nuclear physics

Experiment

- ▶ $(p, p'x)$
 - ▶ RCNP, Japan
 - ▶ iThemba LABS, South Africa
- ▶ $(\gamma, \gamma'x)$, ELI-NP

▶ Oct 2023, RCNP: ^{10}B , ^{11}B , ^{12}C , ^{13}C , ^{27}Al

▶ Oct 2025, RCNP: ^{12}C , ^{27}Al , ^{24}Mg , ^{56}Fe

▶ High-priority to-do: ^6Li , ^7Li , ^9Be (Lightest IVGDR), ^{24}Mg , ^{28}Si , ^{32}S , ^{40}Ca (α -clustering), ^{26}Mg , ^{48}Ca ($N > Z$), ^{14}N , ^{51}V (odd-even and odd-odd effects)

A. Tamii, et al.: *Eur. Phys. J. A*, 59 (2023) 208

- ▶ ELI-NP instruments ready for nuclear data on electric dipole strengths; cross sections for astrophysics, astroparticle physics, energy applications; fission data; and much more
- ▶ Broad collaborations with scientists in related fields in Europe and worldwide
- ▶ Complementary measurements with charged particle accelerators provides a broad range of probes - photons, electrons, hadrons - in the same campus

Acknowledgements: Various topics in this research has been funded by the ELI-RO program by the Institute of Atomic Physics, Măgurele, Romania, contract number ELI-RO/RDI/2024-002 and ELI-RO/RDI/2024-007, the Romanian Ministry of Research, Innovation and Digitization, research contract PN 23 21 01 06.

Special thanks to the nuclear physics colleagues:

- ▶ Dimiter BALABANSKI Head of "Gamma Driven Experiments Department",
- ▶ Senior staff scientists: Andreas OBERSTEDT, Violeta IANCU, Nadia TSONEVA-LARIONOVA
- ▶ Staff scientists: Paul CONSTANTIN, Mihai CUCIUC, Anukul DHAL, Asli KUSOGLU, Haridas PAI, Dmitry TESTOV, Tatiana TOZAR, Gabriel TURTURICA, Yi XU, Adrian ROTARU
- ▶ Postdoctoral staff: Midhun CHERUMUKKU, Sangeeta DHURI, andey JYOTI, Kabita KUNDALIA, Vincent Florian LELASSEUX, Teodora MADGEARU (PETRUSE), Alexandru STATE
- ▶ PhD students: Sara Rebeca BAN, Emanuela BOICU, Andrei COVALI, Andreea GHITIU, Ioana KUNC SER, Cosmina NEDEL CU, Teodora SEBE,
- ▶ Engineers: Sohichiroh AOGAKI, Traian ROMAN