







Coherently enhanced decoherence

Clara Murgui (CERN)

Why quantum sensors?








e . g .




$$\mathcal{L} = \mathcal{L}^{d \leq 4} + \sum_{d=5}^{\infty} \frac{\mathcal{O}_d}{\Lambda_{NP}^{d-4}}$$





e . g . DM



e.g. DM



e.g. DM





“Traditional” direct detection


Nuclear recoils

Lee-Weinberg bound
[Lee, Weinberg,77]


Unitary bound
[Griest, Kamionkowski,90]



WIMPs [2 x miracle]



$$E_{\text{rec}}^{\text{max}} = \frac{q_{\text{max}}^2}{2(A \text{ GeV})} > \text{keV} \quad \text{✂}$$




“Traditional” direct detection


Nuclear recoils

Lee-Weinberg bound
[Lee, Weinberg,77]


Unitary bound
[Griest, Kamionkowski,90]






WIMPs [2 x miracle]




$$E_{\text{rec}}^{\text{max}} = \frac{4m_\chi^2 v_\chi^2}{2(A \text{ GeV})} > \text{keV} \quad \text{✂}$$









fuzzy DM




axions






sterile ν



hidden sector



WIMPs





Also SM!




DEMAND FOR PRECISION
IN LOW ENERGY
PHYSICS (as well!!)

Lowering the threshold

There is plenty of room at the bottom!




WIMPs




How do we “direct-detect”
elusive particles?

Lowering the threshold

Direct detection




Nuclear recoil
PTM exp.



...


~keV energy
resolution




LZ, PandaX,
XENON,
DarkSide,
DEAP...

Lowering the threshold

Direct detection




Nuclear recoil
PTM exp.




...

~keV energy
resolution



~eV energy
resolution


Migdal Effects
Semiconductors




LZ, PandaX,
XENON,
DarkSide,
DEAP...

Lowering the threshold

Direct detection




Nuclear recoil
PTM exp.





...

~keV energy
resolution



~eV energy
resolution



Electron recoils
Migdal Effects
Semiconductors



LZ, PandaX,
XENON,
DarkSide,
DEAP...

Lowering the threshold


Direct detection



Collective excitations


~meV energy resolution

Nuclear recoil
PTM exp.





...

~keV energy resolution



~eV energy resolution



Electron recoils
Migdal Effects
Semiconductors



LZ, PandaX,
XENON,
DarkSide,
DEAP...


Lowering the threshold

Direct detection



Atom
interferometers


nearly
thresholdless



Collective
excitations


~meV energy
resolution

Nuclear recoil
PTM exp.





...

~keV energy
resolution




Electron recoils
Migdal Effects
Semiconductors



LZ, PandaX,
XENON,
DarkSide,
DEAP...


Advantages of low threshold detectors

1 coherent enhancements



Atom
interferometers

nearly
thresholdless





For the i th atom,

$$A_0 e^{iq(x_i-x_0)}$$

Advantages of low threshold detectors

1 coherent enhancements


$$\Gamma \propto \int_{\text{MeV}}^{q_{\text{max}}} dq^2 (\dots)$$




Atom
interferometers


nearly
thresholdless

$$\Gamma = \left| \sum_{i=1}^N A_i \right|^2 \simeq N |A_0|^2$$




Advantages of low threshold detectors

1 coherent enhancements

$$\Gamma \propto \int_0^{q_{\max}} dq^2 (\dots)$$



The graph shows the differential rate $\frac{d\Gamma}{dq^2}$ as a function of q . The curve has a sharp dip at $q = 1/r_c$, which is marked on the x-axis. The area under the curve is shaded in orange.



Atom
interferometers


nearly
thresholdless


$$\Gamma = \left| \sum_{i=1}^N A_i \right|^2 = N^2 |A_0|^2$$



Advantages of low threshold detectors

1 coherent enhancements

$$\Gamma \propto \int_0^{q_{\max}} dq^2 (\dots)$$





Atom
interferometers


nearly
thresholdless

$$\Gamma = \left| \sum_{i=1}^N A_i \right|^2 = N^2 |A_0|^2$$

2 boosted rates

$$\Gamma \propto \int_{q_{\text{th}}}^{q_{\max}} dq^2 \frac{1}{(q^2 + m_{\text{med}}^2)^2} \sim \frac{1}{q_{\text{th}}^2}$$





How do atom interferometers access Γ ?

Atom interferometry

1 atom




(analogy with light)

state preparation

evolution

measurement



Atom interferometry

1 atom




(analogy with light)

state preparation

evolution

measurement



$$|\Delta \mathbf{k}| = \Delta \omega$$

$$\Rightarrow |\Psi(t_p)\rangle = c_g(t_p)|g\rangle \otimes |0\rangle + c_e(t_p)|e\rangle \otimes |k_\gamma\rangle$$

Atom interferometry

1 atom




(analogy with light)

state preparation

evolution

measurement




$$|\Delta \mathbf{k}| = \Delta \omega$$

$$\Rightarrow |\Psi(t_p)\rangle = \cos\left(\frac{\Omega t_p}{2}\right) |g\rangle \otimes |0\rangle + \sin\left(\frac{\Omega t_p}{2}\right) e^{i\phi} |e\rangle \otimes |k_\gamma\rangle$$

Atom interferometry

1 atom




(analogy with light)

state preparation

evolution

measurement





$$|\Delta \mathbf{k}| = \Delta \omega$$

$$\Rightarrow |\Psi(t_{\pi/2})\rangle = \frac{1}{\sqrt{2}} (|g\rangle \otimes |0\rangle + e^{i\phi} |e\rangle \otimes |k_\gamma\rangle)$$

Atom interferometry

1 atom




(analogy with light)

state preparation

evolution

measurement





$$|\Delta \mathbf{k}| = \Delta \omega$$

$$\Rightarrow |\Psi(t_{\pi/2})\rangle = \frac{1}{\sqrt{2}} (|g\rangle \otimes |0\rangle + e^{i\phi} |e\rangle \otimes |k_\gamma\rangle)$$



Atom interferometry

1 atom




(analogy with light)

state preparation

evolution

measurement





$$|\Delta \mathbf{k}| = \Delta \omega$$

$$\Rightarrow |\Psi(t_{\pi/2})\rangle = \frac{1}{\sqrt{2}} (|g\rangle \otimes |0\rangle + e^{i\phi} |e\rangle \otimes |k_\gamma\rangle)$$



Atom interferometry

1 atom




(analogy with light)


state preparation

evolution

measurement





$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$




Atom interferometry

1 atom




(analogy with light)

state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$





measurement

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$


Atom interferometry

1 atom




(analogy with light)

state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$





measurement


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$





measurement

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$


Atom interferometry

1 atom




(analogy with light)

state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$





measurement


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}_{(0)}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}} = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}_{(0)}$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}(\pi/2)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}} (\pi/2) = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}(\pi/2)$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}(\pi/2)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}} = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}(\pi/2)$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}(\pi/2)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\pi/2) = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}(\pi/2)$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}(\pi/2)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}} = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}(\pi/2)$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}(\pi/2)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\pi/2) = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}(\pi/2)$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\{n_g^{(1)}, n_g^{(2)}, \dots, n_g^{(N)}\}(\phi)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{N} \sum_{a=1}^N n_g^{(a)}(\phi)$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement



$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{exp}} (\phi) = \frac{1}{2} (1 + V \cos(\phi + \Delta\phi))$$

The equation shows the experimental result of the measurement. The vertical axis is labeled $\frac{N_g}{N_g + N_e}$ and the horizontal axis is labeled ϕ . The curve is a cosine wave oscillating between 0 and 1. The parameters V and $\Delta\phi$ are highlighted with blue and yellow circles, respectively, and arrows point to them.


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement

$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \text{Tr}\{\rho(\mathcal{T})|g\rangle\langle g|\}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2}(1 + V \cos(\phi + \Delta\phi))$$



V

$\Delta\phi$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{th}}(\phi) = \frac{1}{2}(1 + \cos \phi)$$



$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{exp}}(\phi) = \frac{1}{2}(1 + V \cos(\phi + \Delta\phi))$$

$\uparrow$ $\uparrow$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i(\phi + \Delta\phi)} \\ e^{-i(\phi + \Delta\phi)} & 1 \end{pmatrix}$$

Atom interferometry

1 atom




state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$



measurement

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{th}}(\phi) = \frac{1}{2} (1 + \cos[\phi + \Delta\phi])$$

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{exp}}(\phi) = \frac{1}{2} (1 + V \cos(\phi + \Delta\phi))$$

5th forces


[Wacker, 2010], [Rosi, Sorrentino, et al. 2014]
 [Biedermann, Wu, et al. 2015] [Rosi, D'Amico,
 et al. 2017] [Fray, Diez, et al. 2004] [Schlippert,
 Hartwig, et al. 2014] [Zhou, Long, et al. 2015]
 [Barrett, Antoni-Micollier, et al. 2016] [Kuhn,
 McDonald, et al. 2014] [Barrett, Antoni-
 Micollier, et al. 2015] [Tarallo, Mazzoni, et al
 2014] [Bonnin, Zahzam et al. 2013] [Hartwig,
 Abend, et al. 2015] [Asenbaum, Overstreet, et al
 2020] [Williams, Chiow, et al. 2016] [Battelier,
 Berge, et al., 2019] ...

[Graham, Kaplan, et al. 2016]
 [Arvanitaki, Graham, et al. 2018]
 [Kolb, Weers, et al. 2018]
 [Antypas, Banerjee, 2022]
 [Badurnina, Gipson, et al. 2022]
 [Badurnina, Beniwal, et al. 2023]
 ...

ULDM

[Wicht et al, 2002] [Bennet et
 al. 2006] [Cadoret et al. 2008]
 [Terranova, Tino, 2014]...

EDMs





[Dimopoulos, Graham, et al. 2008] [Hogan, Johnson, et al .
 2011], [Yu, Tinto, 2011] [Graham, Hogan, 2013], [Canuel,
 Bertoldi, et al. 2018] [Canuel, Abend, et al. 2020] [Kolkowitz,
 Pikovski, et al., 2016] [Zhan, Wang, et al. 2020] [El-Neaj,
 Alpigiani, et al. 2020] [Badurina, Bentine, et. Al. 2020]
 [Graham, Hogan, et al. 2016] [Graham, Hogan, et al. 2017],
 [Ballmer, Adhikari, et al. 2022]

GWs


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i(\phi + \Delta\phi)} \\ e^{-i(\phi + \Delta\phi)} & 1 \end{pmatrix}$$

Atom interferometry

1 atom



state preparation



evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$

$$\boxed{\pi/2} + \boxed{\pi} + \boxed{\pi/2}$$

$\underbrace{\hspace{1cm}}_{\mathcal{T}_{1/2}} \quad \underbrace{\hspace{1cm}}_{\mathcal{T}_{1/2}}$

measurement


$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \frac{1}{2} (1 + \cos[\phi + \Delta\phi])$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2} (1 + V \cos(\phi + \Delta\phi))$$

↑ ↑

Atom interferometry

Examples – earth based



UCB

★ **STANFORD** ^{87}Rb

10-m atomic fountain

$r_{\text{cloud}} \sim 0.2\text{mm}, N \sim 10^8$

$\Delta x = 0.1\text{m}, t_{\text{exp}} = 2\text{s}$

★ **LKB**

★ **ZHAOSHAN** ^{87}Rb

12-m atomic fountain



Article

Determination of the fine-structure constant with an accuracy of 81 parts per trillion

<https://doi.org/10.1038/s41586-020-2964-7>

Léo Morel¹, Zhibin Yao¹, Pierre Cladé¹ & Saïda Guellati-Khélifa^{1,2}✉

Received: 7 May 2020



UCB

STANFORD ⁸⁷Rb

10-m atomic fountain

$r_{\text{cloud}} \sim 0.2\text{mm}$, $N \sim 10^8$

$\Delta x = 0.1\text{m}$, $t_{\text{exp}} = 2\text{s}$

EKB

ZHAOSHAN ⁸⁷Rb

12-m atomic fountain

Article

Determination of the fine-structure constant with an accuracy of 81 parts per trillion

<https://doi.org/10.1038/s41586-020-2964-7>

Léo Morel¹, Zhibin Yao¹, Pierre Cladé¹ & Saïda Guellati-Khélifa^{1,2}✉

Received: 7 May 2020

RESEARCH

METROLOGY

Measurement of the fine-structure constant as a test of the Standard Model

Richard H. Parker,^{1*} Chenghui Yu,^{1*} Weicheng Zhong,¹ Brian Estey,¹ Holger Müller^{1,2†}

UCB

STANFORD ⁸⁷Rb

10-m atomic fountain

$$r_{\text{cloud}} \sim 0.2\text{mm}, N \sim 10^8$$

$$\Delta x = 0.1\text{m}, t_{\text{exp}} = 2\text{s}$$

EKB

ZHAOSHAN ⁸⁷Rb

12-m atomic fountain

Article

Determination of the fine-structure constant with an accuracy of 81 parts per trillion

<https://doi.org/10.1038/s41586-020-2964-7>

Léo Morel¹, Zhibin Yao¹, Pierre Cladé¹ & Saïda Guellati-Khélifa^{1,2}✉

Received: 7 May 2020

RESEARCH

METROLOGY

Measurement of the fine-structure constant as a test of the Standard Model

Richard H. Parker,^{1*} Chenghui Yu,^{1*} Weicheng Zhong,¹ Brian Estey,¹ Holger Müll

Test of Equivalence Principle at 10^{-8} Level by a Dual-species Double-diffraction Raman Atom Interferometer

Lin Zhou,^{1,2} Shitong Long,^{1,2,3} Biao Tang,^{1,2} Xi Chen,^{1,2} Fen Gao,^{1,2} Wencui Peng,^{1,2} Weitao Duan,^{1,2,3} Jiaqi Zhong,^{1,2} Zongyuan Xiong,^{1,2} Jin Wang,^{1,2,*} Yuanzhong Zhang,⁴ and Mingsheng Zhan^{1,2,†}

¹State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China

²Center for Cold Atom Physics, Chinese Academy of Sciences, Wuhan 430071, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

(Dated: March 3, 2015)

We report an improved test of the weak equivalence principle by using a simultaneous ^{85}Rb - ^{87}Rb dual-species atom interferometer. We propose and implement a four-wave double-diffraction Raman transition scheme for the interferometer, and demonstrate its ability in suppressing common-mode phase noise of Raman lasers after their frequencies and intensity ratios are optimized. The statistical uncertainty of the experimental data for Eötvös parameter η is 0.8×10^{-8} at 3200 s. With various systematic errors corrected the final value is $\eta = (2.8 \pm 3.0) \times 10^{-8}$. The major uncertainty is attributed to the Coriolis effect.

UCB

STANFORD ^{87}Rb

10-m atomic fountain

$r_{\text{cloud}} \sim 0.2\text{mm}$, $N \sim$
 $\Delta x = 0.1\text{m}$, $t_{\text{exp}} = 2$

Atom-interferometric test of the equivalence principle at the 10^{-12} level

Peter Asenbaum,^{*} Chris Overstreet,^{*} Minjeong Kim, Joseph Curti, and Mark A. Kasevich[†]
Department of Physics, Stanford University, Stanford, California 94305

(Dated: June 25, 2020)

Does gravity influence local measurements? We use a dual-species atom interferometer with 2 s of free-fall time to measure the relative acceleration between ^{85}Rb and ^{87}Rb wave packets in the Earth's gravitational field. Systematic errors arising from kinematic differences between the isotopes are suppressed by calibrating the angles and frequencies of the interferometry beams. We find an Eötvös parameter of $\eta = [1.6 \pm 1.8 \text{ (stat)} \pm 3.4 \text{ (sys)}] \times 10^{-12}$, consistent with zero violation of the equivalence principle. With a resolution of up to $1.4 \times 10^{-11} g$ per shot, we demonstrate a sensitivity to η of $5.4 \times 10^{-11} / \sqrt{\text{Hz}}$.

LKB

ZHAOSHAN ^{87}Rb

12-m atomic fountain

Atom interferometry

Examples – earth based



MAGIS-100

AION

LKB

STANFORD ^{87}Rb

10-m atomic fountain

^{87}Rb

UCB

Atom interferometry

Examples – earth based

AION-REPORT/2023-10



Progress towards ultracold Sr for the AION project – sub-microkelvin atoms and an optical-heterodyne diagnostic tool for injection-locked laser diodes

E. Pasatembou,¹ C. F. A. Baynham,¹ O. Buchmüller,^{1, a)} D. Evans,¹ R. Hobson,¹ L. Iannizzotto-Venezze,¹ and A. Josset¹

Department of Physics, Blackett Laboratory, Imperial College, Prince Consort Road, London, SW7 2AZ, U.K.

(*Electronic mail: c.baynham@imperial.ac.uk)

(Dated: 13 October 2023)



MAGIS-100

UCB

STANFORD ⁸⁷Rb

10-m atomic fountain


AION

LKB

⁸⁷Rb

Atom interferometry

Examples – earth based



Atom interferometry


Examples — space missions



Atom interferometry

Examples — space missions

BECCAL



MAGIS-100

UCB

STANFORD ^{87}Rb

10-m atomic fountain

$$r_{\text{cloud}} \sim 0.2\text{mm}, N \sim 10^8$$
$$\Delta x = 0.1\text{m}, t_{\text{exp}} = 2\text{s}$$

AION

MIGA

ELGAR

LKB

ZAIGA

ZHAOSHAN ^{87}Rb

12-m atomic fountain


Atom interferometry

Examples — space missions

MAQRO SiO_2



BECCAL



GDM

AEDGE

ZAIGA

ZHAOSHAN ^{87}Rb
12-m atomic fountain

AION

MIGA
ELGAR


LKB

MAGIS-100

UCB

STANFORD ^{87}Rb
10-m atomic fountain

$r_{\text{cloud}} \sim 0.2\text{mm}$, $N \sim 10^8$
 $\Delta x = 0.1\text{m}$, $t_{\text{exp}} = 2\text{s}$






How can we detect particles with them?

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}$$

Atom interferometry

1 atom



measurement

evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}]$$

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{th}}(\phi) = \text{Tr}\{\rho(\mathcal{T})|g\rangle\langle g|\}$$

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{exp}}(\phi) = \frac{1}{2}(1 + V \cos(\phi + \Delta\phi))$$

↑



↑

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

1 atom

[Joss, Zeh, 1985]
[Gallis, Fleming, 1990]
[Hornberger, Sipe, 2003]



! $\tau_{1/2}$

$$S(|\mathbf{x}\rangle \otimes |\mathbf{k}\rangle) = |\mathbf{x}\rangle \otimes S_{\{\mathbf{x}\}}|\mathbf{k}\rangle$$



measurement

evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \text{Tr}\{\rho(\mathcal{T})|g\rangle\langle g|\}$$


$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2}(1 + \underset{\uparrow}{V} \cos(\phi + \underset{\uparrow}{\Delta\phi}))$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

1 atom


[Joss, Zeh, 1985]
[Gallis, Fleming, 1990]
[Hornberger, Sipe, 2003]



$$\ln \gamma = - \int_t \int_0^{q_{\max}} dq^2 \frac{d\Gamma}{dq^2} \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

Decoherence Kernel

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = 1 - \exp(i\mathbf{q} \cdot \Delta\mathbf{x})$$



measurement

evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \text{Tr}\{\rho(\mathcal{T})|g\rangle\langle g|\}$$



$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2}(1 + V \cos(\phi + \Delta\phi))$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

1 atom

[Joss, Zeh, 1985]
[Gallis, Fleming, 1990]
[Hornberger, Sipe, 2003]



$$\ln \gamma = - \int_t \int_0^{q_{\max}} dq^2 \frac{d\Gamma}{dq^2} \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

Decoherence Kernel

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = 1 - \exp(i\mathbf{q} \cdot \Delta\mathbf{x})$$



measurement

evolution

$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \frac{1}{2} \left(1 + e^{-\int_q \Gamma(\mathbf{q})(1 - \cos[\mathbf{q} \cdot \Delta\mathbf{x}])} \cos\left[\phi + \int_q \Gamma(\mathbf{q}) \sin[\mathbf{q} \cdot \Delta\mathbf{x}]\right] \right)$$

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2} \left(1 + V \cos(\phi + \Delta\phi) \right)$$

V

$\Delta\phi$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

1 atom

[Joss, Zeh, 1985]
[Gallis, Fleming, 1990]
[Hornberger, Sipe, 2003]


1 $\Gamma \rightarrow 0$



$$\ln \gamma = - \int_t \int_0^{q_{\max}} dq^2 \frac{d\Gamma}{dq^2} \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

Decoherence Kernel

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = 1 - \exp(i\mathbf{q} \cdot \Delta\mathbf{x})$$



measurement

evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \frac{1}{2} \left(1 + \cancel{e^{-\int_q \Gamma(\mathbf{q}) (1 - \cos[\mathbf{q} \cdot \Delta\mathbf{x}])}} \cos[\phi + \cancel{\int_q \Gamma(\mathbf{q}) \sin[\mathbf{q} \cdot \Delta\mathbf{x}]] \right)$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2} \left(1 + \underset{\uparrow}{V} \cos \left(\phi + \underset{\uparrow}{\Delta\phi} \right) \right)$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

1 atom


[Joss, Zeh, 1985]

[Gallis, Fleming, 1990]

[Hornberger, Sipe, 2003]


1 $\Gamma \rightarrow 0$

2 $\Gamma \neq 0, q \ll 1/\Delta x$



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

$$\frac{N_g}{N_g + N_e}$$



tunable phase

Decoherence Kernel

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = 1 - \exp(i\mathbf{q} \cdot \Delta \mathbf{x})$$



measurement

evolution

$$\frac{N_g}{N_g + N_e} \Big|_{\text{th}}(\phi) = \frac{1}{2} \left(1 + e^{-\int_q \Gamma(\mathbf{q}) (1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}])} \cos[\phi + \int_q \Gamma(\mathbf{q}) \sin[\mathbf{q} \cdot \Delta \mathbf{x}]] \right)$$

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\frac{N_g}{N_g + N_e} \Big|_{\text{exp}}(\phi) = \frac{1}{2} (1 + V \cos(\phi + \Delta\phi))$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

1 atom

[Joss, Zeh, 1985]


[Gallis, Fleming, 1990]

[Hornberger, Sipe, 2003]

1 $\Gamma \rightarrow 0$

2 $\Gamma \neq 0, q \ll 1/\Delta x$


3 $\Gamma \neq 0, q \gg 1/\Delta x$



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

Decoherence Kernel

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = 1 - \exp(i\mathbf{q} \cdot \Delta \mathbf{x})$$



measurement

evolution

$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{th}}(\phi) = \frac{1}{2} \left(1 + e^{-\int_q \Gamma(\mathbf{q}) (1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}])} \cos[\phi + \int_q \Gamma(\mathbf{q}) \sin[\mathbf{q} \cdot \Delta \mathbf{x}]] \right)$$

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{exp}}(\phi) = \frac{1}{2} (1 + V \cos(\phi + \Delta\phi))$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$


Atom interferometry

1 atom


[Joss, Zeh, 1985]

[Gallis, Fleming, 1990]

[Hornberger, Sipe, 2003]



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$



Decoherence Kernel

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = 1 - \exp(i\mathbf{q} \cdot \Delta \mathbf{x})$$

evolution

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{th}}(\phi) = \frac{1}{2} (1 + e^{-\int_q \Gamma(\mathbf{q})(1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}])} \cos[\phi + \int_q \Gamma(\mathbf{q}) \sin[\mathbf{q} \cdot \Delta \mathbf{x}])$$


$$\frac{d\hat{\rho}}{dt} = -i[\hat{H}_0 + \hat{H}_{\text{int}}, \hat{\rho}] + \hat{T}\hat{\rho}\hat{T}^\dagger - \frac{1}{2}\{\hat{T}^\dagger\hat{T}, \hat{\rho}\}$$

$$\left. \frac{N_g}{N_g + N_e} \right|_{\text{exp}}(\phi) = \frac{1}{2} (1 + V \cos(\phi + \Delta\phi))$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

N atoms



Statistics


$$\delta\phi|_{\text{shot}} = 1/\sqrt{N}$$

$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

N atoms



?
≡




$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \gamma e^{i\phi} \end{pmatrix}$$

Atom interferometry



N atoms

[Badurina, CM, Plestid, 2024]



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

$\equiv$




$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$


Atom interferometry

[Badurina, CM, Plestid, 2024]

N atoms




?
≡



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

$$|\Psi\rangle = \frac{1}{2^{N/2}} \prod_{i=1}^N (|g\rangle + e^{i\phi} |e\rangle)$$




$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \gamma \end{pmatrix}$$


Atom interferometry

[Badurina, CM, Plestid, 2024]


N atoms



$\equiv$



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$



$$\begin{aligned} |\Psi\rangle &= \frac{1}{2^{N/2}} \prod_{i=1}^N (|g\rangle + e^{i\phi} |e\rangle) \\ &= \frac{1}{2^{N/2}} \sum_{m=0}^N \binom{N}{m} |g_1 \cdots g_m e_{m+1} \cdots e_N\rangle \end{aligned}$$


$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \gamma e^{i\phi} \end{pmatrix}$$

Atom interferometry

N atoms

[Badurina, CM, Plestid, 2024]

[CM, Plestid, 2025]



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$

Decoherence Kernel 1-body

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = G(\mathbf{q}) (1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}] - iN \sin[\mathbf{q} \cdot \Delta \mathbf{x}]) \\ + (1 - G(\mathbf{q})) (1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}] - i \sin[\mathbf{q} \cdot \Delta \mathbf{x}])$$



$$\rho(\mathcal{T}) = \frac{1}{2} \begin{pmatrix} 1 & \gamma e^{i\phi} \\ \gamma^* e^{-i\phi} & \end{pmatrix}$$

Atom interferometry

N atoms

[Badurina, CM, Plestid, 2024]


[CM, Plestid, 2025]



$$\ln \gamma = - \int_{q,t} R(\mathbf{q}) \mathcal{F}_{\text{decoh}}(\mathbf{q})$$


Decoherence Kernel 1-body

$$\mathcal{F}_{\text{decoh}}(\mathbf{q}) = G(\mathbf{q}) (1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}] - iN \sin[\mathbf{q} \cdot \Delta \mathbf{x}]) + (1 - G(\mathbf{q})) (1 - \cos[\mathbf{q} \cdot \Delta \mathbf{x}] - i \sin[\mathbf{q} \cdot \Delta \mathbf{x}])$$



The Debye-Waller factor (cloud form factor squared)

$$G(\mathbf{q}) = \int_y \varrho_y(\{\mathbf{y}\}, \{\mathbf{y}\}) e^{i\mathbf{q} \cdot (\mathbf{y}_i - \mathbf{y}_j)} = e^{-q^2 r_c^2 / 2}$$




$$\Delta\phi = \phi + \gamma_0 + N\gamma$$

How do we test the “coherently enhanced decoherence” at the lab?


Enhanced phase-shift

How do we test it?




Orange square: $\Delta x, N, A, \mathcal{T}, r_c$

Grey square: $\frac{d\sigma}{dq^2}, \frac{d\Phi}{dE}$



Enhanced phase-shift


How do we test it?



Orange square: $\Delta x, N, A, \mathcal{T}, r_c$


Grey square: $\frac{d\sigma}{dq^2}, \frac{d\Phi}{dE}$

$$\Delta\phi = \underbrace{\phi}_{\text{tunable phase}} + \gamma_0 + N\gamma$$



Enhanced phase-shift

How do we test it?



Orange box: $\Delta x, N, A, \mathcal{T}, r_c$

Grey box: $\frac{d\sigma}{dq^2}, \frac{d\Phi}{dE}$


$$\Delta\phi = \underbrace{\phi}_{\text{tunable phase}} + \gamma_0 + N\gamma$$

Incoherent-decoherence

$$\gamma_0 = \int_q \Gamma(\mathbf{q}) [1 - G(\mathbf{q})] \sin(\mathbf{q} \cdot \Delta\mathbf{x})$$

Coherent-decoherence


$$\gamma = \int_q \Gamma(\mathbf{q}) G(\mathbf{q}) \sin(\mathbf{q} \cdot \Delta\mathbf{x})$$



Enhanced phase-shift

How do we test it?

e.g. Kastler Brossel Lab



Orange box: $\Delta x, N, A, \mathcal{T}, r_c$

Grey box: $\frac{d\sigma}{dq^2}, \frac{d\Phi}{dE}$



$$\Delta\phi = \underbrace{\phi}_{\text{tunable phase}} + \gamma_0 + N\gamma$$

Incoherent-decoherence

$$\gamma_0 = \int_q \Gamma(\mathbf{q}) [1 - G(\mathbf{q})] \sin(\mathbf{q} \cdot \Delta\mathbf{x})$$

Coherent-decoherence


$$\gamma = \int_q \Gamma(\mathbf{q}) G(\mathbf{q}) \sin(\mathbf{q} \cdot \Delta\mathbf{x})$$



Enhanced phase-shift @ the lab

[CM, Plestid, 2025]

Via an IR-lamp




Orange square: $\Delta x, N, A, \mathcal{T}, r_c$

Grey square: $\left. \frac{d\sigma}{dq^2} \right|_{\text{IR-lamp}} = \pi \alpha_{\text{pol}}^2 E^2 \left(1 + \left[1 - \frac{q^2}{2E^2} \right]^2 \right)$

Enhanced phase-shift @ the lab

[CM, Plestid, 2025]


Via an IR-lamp



Enhanced phase-shift @ the lab


[CM, Plestid, 2025]

Via an IR-lamp



Orange box: $\Delta x, N, A, \mathcal{T}, r_c$

Black box: $\left. \frac{d\sigma}{dq^2} \right|_{\text{IR-lamp}} = \pi \alpha_{\text{pol}}^2 E^2 \left(1 + \left[1 - \frac{q^2}{2E^2} \right]^2 \right)$



10x10 1mm² LED IR lamps
1€ each

$$\delta\phi(N) = \sqrt{\frac{\delta\phi_{\text{shot}}^2}{N} + \delta\phi_{\text{sys}}^2}$$

(within a single shot)

$$N\gamma \sim 8 \times 10^{-6} \left(\frac{P_{\text{LED}}}{6 \text{ W/mm}^2} \right) \left(\frac{\mathcal{T}_{1/2}}{0.1 \text{ s}} \right) \left(\frac{\Delta x_z}{1 \text{ cm}} \right) \left(\frac{850 \text{ nm}}{\lambda} \right) \left(\frac{N}{10^6} \right) \left(\frac{\alpha_{\text{pol}}}{47.39^\circ \text{Å}^3} \right)^2$$

[CM, Plestid, 2025]

Lindblad eq



$$\blacksquare \quad \left. \frac{d\sigma}{dq^2} \right|_{e^- \text{ gun}} = 4\pi Z^2 r_A^4 \alpha_{\text{EM}}^2 \frac{1}{\beta_e^2}$$



$$N\gamma \simeq 6 \times 10^{-3} \left(\frac{\mathcal{T}_{1/2}}{0.1 \text{ s}} \right) \left(\frac{\Delta x}{1 \text{ m}} \right) \left(\frac{I_e}{1 \text{ A}} \right) \left(\frac{0.1 \text{ mm}}{r_c} \right)^6 \left(\frac{10^6}{N} \right)$$

$$\gamma_0 \simeq 5 \times 10^{-4} \left(\frac{\mathcal{T}_{1/2}}{0.1 \text{ s}} \right) \left(\frac{1 \text{ m}}{\Delta x} \right) \left(\frac{I_e}{1 \text{ A}} \right) \left(\frac{0.1 \text{ mm}}{r_c} \right)^2$$

https://www.kimballphysics.com/learning_center/electron-gun-beam-systems/


Dark matter inprints

[Riedel, 2013]

[Riedel, Yavin, 2017]

[Du, CM, Pardo, Wang, Zurek, 2022]

[CM, Plestid, 2025]




Orange square: $\Delta x, N, A, \mathcal{T}, r_c$

Dark grey square: $\left. \frac{d\sigma}{dq^2} \right|_{\text{DM}} = 4\pi A^2 \alpha_{\text{DM}}^2 \frac{1}{u_\chi^2} \left(\frac{1}{q^2 + m_\phi^2} \right)^2$

coherent decoherence dominates

$$\alpha_{\text{DM}} < 10^{-16} \left(\frac{m_\chi}{10 \text{ eV}} \right) \left(\frac{1 \text{ cm}}{\Delta x} \right)^{\frac{1}{2}} \left(\frac{87}{A} \right) \\ \times \left(\frac{0.1 \text{ s}}{\mathcal{T}_{1/2}} \right)^{\frac{1}{2}} \left(\frac{10^6}{N} \right)^{\frac{3}{4}} \left(\frac{1}{N} \right)^{\frac{1}{4}} \left(\frac{1}{f_\chi} \right)^{1/2} 10^2 \left[\left(\frac{m_\phi r_c}{10} \right)^2 \right]$$



$$\Rightarrow \alpha_{\text{DM}} = y_\chi y_N / 4\pi$$


Dark matter inprints

[Riedel, 2013]

[Riedel, Yavin, 2017]


[Du, **CM**, Pardo, Wang, Zurek, 2022]

[**CM**, Plestid, 2025]



coherent decoherence dominates

$$\alpha_{\text{DM}} < 10^{-16} \left(\frac{m_\chi}{10 \text{ eV}} \right) \left(\frac{1 \text{ cm}}{\Delta x} \right)^{\frac{1}{2}} \left(\frac{87}{A} \right) \\ \times \left(\frac{0.1 \text{ s}}{\mathcal{T}_{1/2}} \right)^{\frac{1}{2}} \left(\frac{10^6}{N} \right)^{\frac{3}{4}} \left(\frac{1}{N} \right)^{\frac{1}{4}} \left(\frac{1}{f_\chi} \right)^{1/2} 10^2 \left[\left(\frac{m_\phi r_c}{10} \right)^2 \right]$$




$$\Rightarrow \alpha_{\text{DM}} = y_\chi y_N / 4\pi$$

Future steps

Explore new observables

[[CM](#), [Plestid](#), 251X.XXXX]



Fluctuations in atom interferometers as a new tool for dark matter

Clara Murgui^{1,*} and Ryan Plestid^{1,†}

¹*Theoretical Physics Department, CERN, 1 Esplanade des Particules, CH-1211 Geneva 23, Switzerland*

(Dated: November 18, 2025)

Disadvantages of coherent enhancements

Penalty if the “most preferred” incoming momentum is $\neq \mathcal{O}(1/r_c)$

$$\text{obs} \propto \int dt \int dE \frac{d\Phi}{dE} \int dq^2 \frac{d\sigma}{dq^2} G(\mathbf{q}, t) \int \frac{d\phi}{(2\pi)^3} \sin(\mathbf{q} \cdot \mathbf{v})$$

Not applicable to spin-dependent interactions (γ_5)



The weirdos: coherent states

[Badurina, CM, Plestid, 24] [CM, Plestid, 251X.XXXX]

The mean

$$\langle \mathcal{O}_+ \rangle = \text{Tr}\{\rho \mathcal{O}_+\} = N \text{Tr}\{\rho_1 |+\rangle \langle +|\} = \frac{N}{2} (1 + \text{Re}[\blacksquare_1])$$

$$\langle n_+ \rangle = \frac{1}{N} \sum_{i=1}^N n_+^{(a)}$$

$$\rho_1 = \frac{1}{2} \begin{pmatrix} \circ & \blacksquare \\ \blacksquare & \circ \end{pmatrix}$$

Pure product state

$$\circ = 1 \quad \blacksquare = e^{i\Phi}$$

$$\langle O_+ \rangle|_{\text{pure}} \propto N$$

Fully decohered $e^{-(s \nearrow s_0)^0}$

$$\circ = 1 \quad \blacksquare = 0$$

$$\langle O_+ \rangle|_{\text{decoh}} \propto N$$

The weirdos: coherent states

[Badurina, CM, Plestid, 24] [CM, Plestid, 251X.XXXX]

The mean

$$\langle \mathcal{O}_+ \rangle = \text{Tr}\{\rho \mathcal{O}_+\} = N \text{Tr}\{\rho_1 |+\rangle \langle +|\} = \frac{N}{2} (1 + \text{Re}[\blacksquare_1])$$

Pure product state

$$\circ = 1 \quad \blacksquare = e^{i\Phi}$$

$$\langle \mathcal{O}_+ \rangle|_{\text{pure}} \propto N$$

$$\langle n_+ \rangle = \frac{1}{N} \sum_{i=1}^N n_+^{(a)}$$

$$\rho_1 = \frac{1}{2} \begin{pmatrix} \circ & \blacksquare \\ \blacksquare & \circ \end{pmatrix}$$

Fully decohered $e^{-(s+s_0)}$

$$\circ = 1 \quad \blacksquare = 0$$

$$\langle \mathcal{O}_+ \rangle|_{\text{decoh}} \propto N$$

The variance

$$\cos 2\Phi = 2 \cos^2 \Phi - 1$$

$$\sigma_+^2 = \langle \mathcal{O}_+^2 \rangle - \langle \mathcal{O}_+ \rangle^2 = \frac{N}{8} \left[N (1 + \text{Re}[\star] - 2\text{Re}[\blacksquare_1]^2) + 1 - \text{Re}[\star] \right]$$

Pure product state

$$\circ = 1 \quad \blacksquare = e^{i\Phi} \quad \star = e^{i2\Phi}$$

$$\sigma_+|_{\text{pure}} \propto \sqrt{N}$$

Fully decohered $e^{-(s+s_0)}$

$$\circ = 1 \quad \blacksquare = 0 \quad \star = 0$$

$$\sigma_+|_{\text{decoh}} \propto N$$

$$s = \frac{1}{N} \sum_{a=1}^N [n_+^{(a)} - \langle n_+ \rangle]^2$$

$$\rho_2 = \frac{1}{4} \begin{pmatrix} \circ & \blacksquare & \blacksquare & \star \\ \blacksquare & \circ & \circ & \blacksquare \\ \blacksquare & \circ & \circ & \blacksquare \\ \star & \blacksquare & \blacksquare & \circ \end{pmatrix}$$


Future steps

Explore new observables

[[CM](#), Plestid, 251X.XXXX]

Add more generic configurations to the formalism

⇒ Squeezing



[Treutlein, et al., 18]

Future steps


Explore new observables

[CM, Plestid, 251X.XXXX]


Add more generic configurations to the formalism

⇒ Squeezing

⇒ Liquids



[Simonds et al., 2001, Nature]



Future steps

Explore new observables

[[CM](#), Plestid, 251X.XXXX]

Add more generic configurations to the formalism


⇒ Squeezing

⇒ Liquids

Convince experimentalists!

Conclusions


Atom interferometers open (“THE”) parameter space



Conclusions

Atom interferometers open (“THE”) parameter space

These experiments exist! Many funded already. Parasitic searches.




[AION collaboration, Badurina et al, 2019]

Conclusions

Atom interferometers open (“THE”) parameter space

These experiments exist! Many funded already. Parasitic searches

Ambitious projects require controlled backgrounds (SM physics)



[Du, CM, Pardo, Wang, Zurek, 2023]

Conclusions

Atom interferometers open (“THE”) parameter space

These experiments exist! Many funded already. Parasitic searches

Ambitious projects require controlled backgrounds (SM physics)

Boundaries not reached yet!

The line between science fiction and science fact vanishes quickly

Conclusions


Atom interferometers open (“THE”) parameter space

These experiments exist! Many funded already. Parasitic searches

Ambitious projects require controlled backgrounds (SM physics)

Boundaries not reached yet!

The line between science fiction and science fact vanishes quickly



THANK YOU!

