

Cornering New Physics through precision measurements at FCC-ee

Electroweak observables, rare decays, and other things

Lukas Allwicher

Seminar, IJCLab Orsay, 18.12.2025

Outline

I Context

- Precision measurements as BSM probes
- SMEFT (and flavour assumptions)
- Current status

II FCC-ee

- Electroweak Precision Observables
- Flavour prospects
- Above the Z -pole

Context



New Physics

Several issues with the Standard Model, e.g.

- > Dark Matter
- > Neutrino Masses
- > Flavour Puzzle
- > Hierarchy problem

$$h \text{ --- } \textcircled{X} \text{ --- } h \longrightarrow \delta m_h^2 \approx \frac{g^2}{16\pi^2} M_X^2$$

⇒ Need “light” NP to avoid tuning, **TeV Scale**

All other problems may have a solution at the TeV scale as well (WIMP, ISS, non-universal gauge interactions, ...) → worth exploring!



We are only now starting to probe the TeV scale systematically

What more could we learn from future precision measurements?

Precision measurements as BSM probes


- > Through indirect effects, precision measurements can give insights into scales much higher than the ones involved in the process
- > Particularly powerful when the SM rate for a process is suppressed:
 - FCNCs (e.g. $b \rightarrow s \ell \ell$)
 - LFV (e.g. $\mu \rightarrow e \gamma$)
- > In recent years, also tests of universality
 - LFU ratios ($R_K, R_D, \dots$)

$$\frac{1}{M^2} \leftrightarrow \frac{1}{16\pi^2} \frac{\varepsilon}{v^2}$$



Past examples

The Higgs mass




- > If you miss some piece of information about a model, precision tests can help directing/excluding scenarios

$K - \bar{K}$ mixing

- > CP violation in neutral kaon mixing was a New Physics signal
- > Ultimately leading to the discovery of the third generation
- > Today, e.g. a measurement of LFV would clearly be a BSM signal in the same way

Current status

- > In the absence of any BSM signals, set bounds
- > Each observable mediated by a contact interaction $\frac{1}{\Lambda^2} \mathcal{O}_{\text{eff}}^{d=6} \rightarrow \Lambda > \dots$
- > European Strategy for Particle Physics '26



What is the scale of New Physics?

- > Depends on the assumptions! Doesn't have to be 10^7 TeV ($\leftrightarrow$ anarchic flavour)
- > Take again $K - \bar{K}$ mixing.
After discovery, “superweak” theory


[Wolfenstein '79]

$$\frac{ce^{i\phi}}{\Lambda^2} (\bar{s}\gamma^\mu d)^2 \rightarrow \Lambda \sim 10^4 \text{ TeV}$$

- > In the SM:

$$\frac{c}{\Lambda^2} = \frac{(V_{td}V_{ts}^*)^2 G_F^2 m_t^2}{16\pi^2}$$

What is the flavour structure of low-scale NP?



The New Physics Flavour Puzzle

Exploring the UV systematically

Standard Model Effective Field Theory

- In presence of a mass-gap, somewhat model-independent approach
→ study classes of models
- Complement the SM with a tower of higher-dimensional operators, with SM fields and gauge symmetry

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \frac{1}{\Lambda^2} \sum_i \mathcal{C}_i \mathcal{O}_i^{(6)} + \dots$$

- Correct way of dealing with scale separations (RGE)
- Stop the expansion at $d = 6$
- 59 operator structures, 2499 independent coefficients
- Most of the parameters come from flavour → **flavour assumptions**

- Minimal Flavour Violation: only breaking $U(3)^5$ symmetry comes from SM Yukawas
- e.g. $q_L \sim (\mathbf{3}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{1})$ under

$$U(3)^5 \equiv U(3)_q \times U(3)_\ell \times U(3)_u \times U(3)_d \times U(3)_e$$

- $Y_u \sim (\mathbf{3}, \mathbf{1}, \bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$
- Yukawas promoted to spurions keeping track of $U(3)^5$ breaking

$$\bar{q}_L^i \gamma^\mu q_L^j (a \delta_{ij} + b [Y_u Y_u^\dagger]_{ij} + \dots)$$

- Good to suppress flavour-changing processes:

$$\lambda_{\text{FC}} \approx (Y_u Y_u^\dagger)_{\text{FC}} \sim \begin{pmatrix} 0 & \lambda^5 & \lambda^3 \\ \lambda^5 & 0 & \lambda^2 \\ \lambda^3 & \lambda^2 & 0 \end{pmatrix} \quad \lambda \simeq 0.2$$

- But:** leading term is flavour-conserving and **universal**
→ collider searches push the scale to $\Lambda \gtrsim 10 \text{ TeV}$

Flavour assumptions: $U(2)^5$

- Yukawa terms break the $U(3)^5$ symmetry:

$$U(3)^5 \xrightarrow{\mathcal{L}_{\text{Yukawa}}} U(1)_B \times U(1)_L^3$$

- However, light family Yukawas very small: approximate $U(2)^5$ symmetry

$$Y \simeq y_3 \left(\begin{array}{cc|c} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \hline 0 & 0 & 1 \end{array} \right) \quad [Barbieri, Isidori, Lodone, Straub 1105.2296] \quad U(2)^5 = U(2)_q \times U(2)_\ell \times U(2)_u \times U(2)_d \times U(2)_e$$

e.g.

$$q_L^{i=1,2} \sim (\mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{1}) \quad q_L^3 \text{ singlet}$$

Minimal breaking:

$$Y = y_3 \left(\begin{array}{c|c} \Delta & V \\ \hline 0 & 1 \end{array} \right) \quad \begin{array}{ll} V \sim (\mathbf{2}, \mathbf{1}) & \Delta \sim (\mathbf{2}, \bar{\mathbf{2}}) \\ |V_q| = \mathcal{O}(y_t V_{ts}) & |\Delta| \sim y_{c,s,\mu} \end{array}$$

$U(2)^5$ symmetry at work

- ψ^2 operators: e.g. $\mathcal{C}_{He}$

$$\mathcal{L}_{\text{SMEFT}} \supset [\mathcal{C}_{He}]_{ij} (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{e}_i \gamma^\mu e_j) \\ \xrightarrow{U(2)^5} \mathcal{C}_{He}^{[33]} (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{e}_3 \gamma^\mu e_3) + \mathcal{C}_{He}^{[ii]} (H^\dagger i \overleftrightarrow{D}_\mu H) \sum_{i=1}^2 (\bar{e}_i \gamma^\mu e_i)$$

- $6 \rightarrow 2$ independent coefficients
- ψ^4 operators: e.g. $\mathcal{C}_{lequ}$

$$\mathcal{L}_{\text{SMEFT}} \supset [\mathcal{C}_{lequ}]_{ijkl} (\bar{\ell}_i e_j) (\bar{q}_k u_l) \xrightarrow{U(2)^5} \mathcal{C}_{lequ}^{[3333]} (\bar{\ell}_3 e_3) (\bar{q}_3 u_3)$$

- $81 \rightarrow 1$ independent coefficients!

The $U(2)$ -symmetric SMEFT

- 2499 independent parameters at $d = 6$
- exact $U(2)$: 124 CPC + 23 CPV

[Faroughy, Isidori, Wilsch, Yamamoto 2005.05366]



Operators	$U(2)^5$ [terms summed up to different orders]													
	Exact		$\mathcal{O}(V^1)$		$\mathcal{O}(V^2)$		$\mathcal{O}(V^1, \Delta^1)$		$\mathcal{O}(V^2, \Delta^1)$		$\mathcal{O}(V^2, \Delta^1 V^1)$		$\mathcal{O}(V^3, \Delta^1 V^1)$	
Class 1-4	9	6	9	6	9	6	9	6	9	6	9	6	9	6
$\psi^2 H^3$	3	3	6	6	6	6	9	9	9	9	12	12	12	12
$\psi^2 XH$	8	8	16	16	16	16	24	24	24	24	32	32	32	32
$\psi^2 H^2 D$	15	1	19	5	23	5	19	5	23	5	28	10	28	10
$(\bar{L}L)(\bar{L}L)$	23	–	40	17	67	24	40	17	67	24	67	24	74	31
$(\bar{R}R)(\bar{R}R)$	29	–	29	–	29	–	29	–	29	–	53	24	53	24
$(\bar{L}L)(\bar{R}R)$	32	–	48	16	64	16	53	21	69	21	90	42	90	42
$(\bar{L}R)(\bar{R}L)$	1	1	3	3	4	4	5	5	6	6	10	10	10	10
$(\bar{L}R)(\bar{L}R)$	4	4	12	12	16	16	24	24	28	28	48	48	48	48
total:	124	23	182	81	234	93	212	111	264	123	349	208	356	215

Table 6: Number of independent operators in the SMEFT assuming a minimally broken $U(2)^5$ symmetry, including breaking terms up to $\mathcal{O}(V^3, \Delta^1 V^1)$. Notations as in Table 1.


How low can the NP scale be?

- consider collider, electroweak and flavour observables

Data at different energy scales




From D. Hitlin's slides at FermiLab



RGE

Connecting scales



- > Assume heavy NP at scale $\Lambda_{\text{UV}} \gtrsim \text{TeV}$
- > After matching to SMEFT, use the SMEFT RGE to connect to lower scales
- > For flavour observables, need to integrate out h , W , Z , t at the EW scale
- > LEFT RGE to the scale of the observable
- > LL approximation: $\mathcal{C}_i(\mu) = \frac{1}{16\pi^2} \log \frac{\mu}{\Lambda} \gamma_{ij} \mathcal{C}(\Lambda)$

Flavour: $U(2)$ needs to be broken

- > It is broken already at dimension four
- > Use spurions to parametrise the breaking

Exact $U(2)^5$

$$\bar{q}_L^3 \gamma_\mu q_L^3 + \epsilon \bar{q}_L^i \gamma_\mu q_L^i$$

- No flavour violation
- Get bounds from collider and EW observables

Minimally broken $U(2)^5$

$$\bar{q}_L^i V_q^i \gamma_\mu q^3 \quad V_q \sim \mathcal{O} \begin{pmatrix} V_{td} \\ V_{ts} \end{pmatrix}$$

- Flavour violating couplings
- Effects mainly in B physics ($b \rightarrow c, s$ transitions)

In single-parameter fits: **alignment** of the left-handed quark doublet:

down-aligned


$$q_L^{\text{down}} = \begin{pmatrix} V_{\text{CKM}}^\dagger u_L \\ d_L \end{pmatrix}$$

up-aligned

$$q_L^{\text{up}} = \begin{pmatrix} u_L \\ V_{\text{CKM}} d_L \end{pmatrix}$$

SMEFT fits to $U(2)$ operators

- > From 10^6 to ~ 10 TeV through $U(2)^5$
- > Single-coefficient fits



New Physics in the third generation?

- > **Idea:** New Physics couples predominantly to the third generation
- > Mimicks the SM Yukawas ($\leftrightarrow$ SM flavour puzzle?)
- > Couplings to light families dynamically suppressed
- > In the EFT, this is an extra assumption
- > An accidental $U(2)$ symmetry on the light families arises naturally
- > Starting from the $U(2)$ -symmetric case, add extra suppressions "by hand"

$$\bar{q}_L^i \gamma^\mu q_L^j \rightarrow a \bar{q}_L^3 \gamma^\mu q_L^3 + b \sum_{i=1}^2 \bar{q}_L^i \gamma^\mu q_L^i$$


- > a and b independent coefficients
- > E.g. for quarks $q_i \rightarrow \varepsilon_Q q_i$ for $i = 1, 2$ (assume $b \ll a$)

New Physics in the third generation?

[LA, Cornella, Isidori, Stefaneke 2311.00020]

- > Suppress operators with light fermion indices
- > Λ_{NP} still compatible with $\sim$ TeV under non-tuned conditions

■ Flavor ■ EW ■ Collider



Where do we go from here?

- > Constraints from processes involving third-generation fields are the weakest
- > If NP couples mainly to the third gen., it could still very well lie around the TeV scale
- > Interesting not also in view of the hierarchy problem, but also of the rich ongoing and future experimental programme (LHC experiments or Belle-II might still be able to see a NP signal)
- > Especially in the absence of clear NP indications, precision measurements make more sense than ever
- > FCC-ee provides an incredible framework for indirect searches

FCC-ee



Indirect searches at FCC-ee

[Table from FCC FSR 2404.12809]

Working point	Z pole	WW thresh.	ZH	$t\bar{t}$	
$\sqrt{s}$ (GeV)	88, 91, 94	157, 163	240	340–350	365
Lumi/IP ($10^{34} \text{ cm}^{-2} \text{ s}^{-1}$)	140	20	7.5	1.8	1.4
Lumi/year (ab^{-1})	68	9.6	3.6	0.83	0.67
Run time (year)	4	2	3	1	4
Integrated lumi. (ab^{-1})	205	19.2	10.8	0.42	2.70
Number of events	6×10^{12} Z	2.4×10^8 WW	2.2×10^6 ZH + 65k WW $\rightarrow$ H	2×10^6 $t\bar{t}$ + 370k ZH + 92k WW $\rightarrow$ H	

EWPOs

- > Focus on Z - and W -pole obs.
- > Constrain flavour-conserving NP

Flavour

- > Large number of $b\bar{b}$ and $\tau^+\tau^-$ pairs from Z decays
- > Constrain flavour-violating NP

Higgs

- > Yukawas, hVV couplings
- > Self-coupling

→ overall great potential for cornering NP indirectly

EWPOs at FCC-ee

- Measure Z - and W boson couplings to fermions through partial decay widths
- Current measurements largely from LEP, $\mathcal{O}(1\%)$ precision
- Expect an improvement by 10-100 at FCC-ee

	Observable	Definition
Z-pole	Γ_Z	$\sum_f \Gamma(Z \rightarrow f\bar{f})$
	σ_{had}	$\frac{12\pi}{m_Z} \frac{\Gamma(Z \rightarrow e^+e^-)\Gamma(Z \rightarrow q\bar{q})}{\Gamma_Z^2}$
	$R_f (f = e, \mu, \tau, c, b)$	$\frac{\Gamma(Z \rightarrow f\bar{f})}{\sum_q \Gamma(Z \rightarrow q\bar{q})}$
	$A_f (f = e, \mu, \tau, s, c, b)$	$\frac{\Gamma(Z \rightarrow f_L f_L) - \Gamma(Z \rightarrow f_R \bar{f}_R)}{\Gamma(Z \rightarrow f\bar{f})}$
	$A_{\text{FB}}^{0,\ell} (\ell = e, \mu, \tau)$ $A_q^{\text{FB}} (q = c, b)$	$\frac{3}{4} A_e A_\ell$ $\frac{3}{4} A_e A_q$
W-pole	m_W	$\sum_{f_1, f_2} \Gamma(W \rightarrow f_1 f_2)$
	Γ_W $\text{Br}(W \rightarrow \ell\nu) (\ell = e, \mu, \tau)$	

[Table from FCC FSR 2505.00274]

Observable	present value	present $\pm$ uncertainty	FCC-ee Stat.	FCC-ee Syst.	Comment and leading uncertainty
m_Z (keV)	91 187 600	$\pm$ 2000	4	100	From Z line shape scan Beam energy calibration
Γ_Z (keV)	2 495 500	$\pm$ 2300	4	12	From Z line shape scan Beam energy calibration
$\sin^2 \theta_W^{\text{eff}} (\times 10^6)$	231,480	$\pm$ 160	1.2	1.2	From $A_{\text{FB}}^{\mu\mu}$ at Z peak Beam energy calibration
$1/\alpha_{\text{QED}}(m_Z^2) (\times 10^3)$	128 952	$\pm$ 14	3.9 0.8	small thc	From $A_{\text{FB}}^{\mu\mu}$ off peak From $A_{\text{FB}}^{\mu\mu}$ on peak QED&EW uncert. dominate
$R_\ell^Z (\times 10^3)$	20 767	$\pm$ 25	0.05	0.05	Ratio of hadrons to leptons Acceptance for leptons
$\alpha_S(m_Z^2) (\times 10^4)$	1 196	$\pm$ 30	0.1	1	Combined $R_\ell^Z, \Gamma_{\text{tot}}^Z, \sigma_{\text{had}}^0$ fit
$\sigma_{\text{had}}^0 (\times 10^3)$ (nb)	41 480.2	$\pm$ 32.5	0.03	0.8	Peak hadronic cross section Luminosity measurement
$N_\nu (\times 10^3)$	2 996.3	$\pm$ 7.4	0.09	0.12	Z peak cross sections Luminosity measurement
$R_b (\times 10^6)$	216 290	$\pm$ 660	0.25	0.3	Ratio of $b\bar{b}$ to hadrons
$A_{\text{FB}}^{b,0} (\times 10^4)$	992	$\pm$ 16	0.04	0.04	b-quark asymmetry at Z pole From jet charge
$A_{\text{FB}}^{\text{pol},\tau} (\times 10^4)$	1 498	$\pm$ 49	0.07	0.2	τ polarisation asymmetry τ decay physics

EWPOs in SMEFT at tree level

$$\mathcal{C}_{H\psi}(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{\psi}\gamma^\mu\psi) \xrightarrow{H \rightarrow \langle H \rangle} -\frac{v^2}{2}\mathcal{C}_{H\psi}Z_\mu(\bar{\psi}\gamma^\mu\psi)$$

> Sensitive to 23 parameters

$[\mathcal{O}_{H\ell}^{(1)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{\ell}_p \gamma^\mu \ell_r)$	$[\mathcal{O}_{Hu}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$	m_W “S-parameter” G_F
$[\mathcal{O}_{H\ell}^{(3)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{\ell}_p \tau^I \gamma^\mu \ell_r)$	$[\mathcal{O}_{Hd}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$	
$[\mathcal{O}_{Hq}^{(1)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$	$\mathcal{O}_{HD} = H^\dagger D_\mu H ^2$	
$[\mathcal{O}_{Hq}^{(3)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$	$\mathcal{O}_{HWB} = (H^\dagger \tau^I H) W_{\mu\nu}^I B^{\mu\nu}$	
$[\mathcal{O}_{He}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$	$[\mathcal{O}_{\ell\ell}]_{1221} = (\bar{\ell}_1 \gamma^\mu \ell_2)(\bar{\ell}_2 \gamma^\mu \ell_1)$	


> $\mathcal{A} = \mathcal{A}_0 \left(\frac{g_{\text{SM}}}{m_W^2} + \frac{g_{\text{NP}}}{\Lambda_{\text{NP}}^2} \right)$

> 10^5 improvement in statistics: $\Lambda_{\text{NP}} \rightarrow \simeq 20 \Lambda_{\text{NP}}^{\text{now}}$

> Naive scaling, systematics (and theory errors) have an effect at such precision


Projections for tree-level operators

- > $U(2)^5$ symmetry acting on the light generations imposed on SMEFT
- > From current $\mathcal{O}(\text{few})\text{TeV}$ to 30 TeV range at FCC-ee



Sizeable RGE effects


- > Loop suppression $\frac{1}{16\pi^2} g^2 \log \frac{m_Z}{\Lambda} \simeq -(0.14)^2$ for $g = 1$ and $\Lambda = 2 \text{ TeV}$
- > Can still probe loop-suppressed effects up to the few TeV range
- > LL effects:




$$[\dot{C}_{Hl}^{(3)}]_{\alpha\beta} \supset 2N_c [C_{lq}^{(3)}]_{\alpha\beta kl} [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{lk}$$

- Particularly relevant if y_t is involved
- Effects e.g. in $Z \rightarrow \tau\tau$

- > Sometimes even NLL effects can be relevant



$\rightarrow C_{Hu}$




$\rightarrow C_{HD} \rightarrow \delta m_W$

Projections from loop-level EWPOs

[LA, Cornella, Isidori, Stefaneke 2311.00020]

> $\mathcal{O}(10)$ TeV constraints for four-fermion operators (3rd gen. quarks)



From SMEFT to the UV

- > SMEFT as a “bookkeeping” tool
- > Both single-parameter fits and global fits do not carry the same information as an explicit model
- > Not all of the SMEFT parameter space can be spanned by UV models
- > In particular, flat directions may not be populated
- > Even if populated at tree-level, loops (RGE) will typically break them

Which models with heavy NP can be probed at FCC-ee? Which scales?

Linear extensions of the Standard Model

[De Blas et al. 1711.10391]

- > Finite number of new states can couple linearly to the SM fields: *Granada dictionary*
- > Matching to SMEFT at $d = 6$ well known

Scalar	$\mathcal{S}$	$\mathcal{S}_1$	$\mathcal{S}_2$	φ	Ξ	Ξ_1	Θ_1	Θ_3
	$(1, 1)_0$	$(1, 1)_1$	$(1, 1)_2$	$(1, 2)_{\frac{1}{2}}$	$(1, 3)_0$	$(1, 3)_1$	$(1, 4)_{\frac{1}{2}}$	$(1, 4)_{\frac{3}{2}}$
	ω_1	ω_2	ω_4	Π_1	Π_7	ζ		
Fermion	$(3, 1)_{-\frac{1}{3}}$	$(3, 1)_{\frac{2}{3}}$	$(3, 1)_{-\frac{4}{3}}$	$(3, 2)_{\frac{1}{6}}$	$(3, 2)_{\frac{7}{6}}$	$(3, 3)_{-\frac{1}{3}}$		
	Ω_1	Ω_2	Ω_4	Υ	Φ			
	$(6, 1)_{\frac{1}{3}}$	$(6, 1)_{-\frac{2}{3}}$	$(6, 1)_{\frac{4}{3}}$	$(6, 3)_{\frac{1}{3}}$	$(8, 2)_{\frac{1}{2}}$			
Vector	N	E	Δ_1	Δ_3	Σ	Σ_1		
	$(1, 1)_0$	$(1, 1)_{-1}$	$(1, 2)_{-\frac{1}{2}}$	$(1, 2)_{-\frac{3}{2}}$	$(1, 3)_0$	$(1, 3)_{-1}$		
	U	D	Q_1	Q_5	Q_7	T_1	T_2	
	$(3, 1)_{\frac{2}{3}}$	$(3, 1)_{-\frac{1}{3}}$	$(3, 2)_{\frac{1}{6}}$	$(3, 2)_{-\frac{5}{6}}$	$(3, 2)_{\frac{7}{6}}$	$(3, 3)_{-\frac{1}{3}}$	$(3, 3)_{\frac{2}{3}}$	
Vector	$\mathcal{B}$	$\mathcal{B}_1$	$\mathcal{W}$	$\mathcal{W}_1$	$\mathcal{G}$	$\mathcal{G}_1$	$\mathcal{H}$	$\mathcal{L}_1$
	$(1, 1)_0$	$(1, 1)_1$	$(1, 3)_0$	$(1, 3)_1$	$(8, 1)_0$	$(8, 1)_1$	$(8, 3)_0$	$(1, 2)_{\frac{1}{2}}$
	$\mathcal{L}_3$	$\mathcal{U}_2$	$\mathcal{U}_5$	$\mathcal{Q}_1$	$\mathcal{Q}_5$	$\mathcal{X}$	$\mathcal{Y}_1$	$\mathcal{Y}_5$
	$(1, 2)_{-\frac{3}{2}}$	$(3, 1)_{\frac{2}{3}}$	$(3, 1)_{\frac{5}{3}}$	$(3, 2)_{\frac{1}{6}}$	$(3, 2)_{-\frac{5}{6}}$	$(3, 3)_{\frac{2}{3}}$	$(\bar{6}, 2)_{\frac{1}{6}}$	$(\bar{6}, 2)_{-\frac{5}{6}}$

All (except very few) new states are probed by EWPOs at one-loop

Analysis strategy


- > Assume SM central values + FCC-ee expected errors
- > Integrate out one state of the Granada dictionary at a time, assuming couplings are defined at a scale $\Lambda = 2 \text{ TeV}$
- > Consider only couplings at $d \leq 4$
- > RGE evolve from 2 TeV to m_Z (first LL only)
→ use `DsixTools` [2010.16341]
- > Get bounds on the mediator mass for $y, g, \lambda = 1$
(dimensionful couplings $\kappa = 5 \text{ TeV}$)
- > Flavour assumptions:
 - (a) **Flavour-universal** couplings: $y_{11} = y_{22} = y_{33}$ (or $y_1 = y_2 = y_3$ for fermions), and $y_{ij} = 0$ for $i \neq j$.
 - (b) **Third-generation only** couplings: $y_{ij} = y \delta_{i3} \delta_{j3}$

Scalars

(*) = special choice of couplings to avoid tree-level EWPO

■ Universal couplings ■ Third-gen. only ■ Flavourless couplings ■ Antisymm. couplings

▨ = no running



[1711.1039]

Name	S	S_1	S_2	φ	Ξ	Ξ_1	Θ_1	Θ_3
Irrep	$(1,1)_0$	$(1,1)_{\frac{1}{2}}$	$(1,1)_2$	$(1,2)_{\frac{1}{2}}$	$(1,3)_0$	$(1,3)_1$	$(1,4)_{\frac{1}{2}}$	$(1,4)_{\frac{3}{2}}$

Name	ω_1	ω_2	ω_4	Π_1	Π_7	ζ
Irrep	$(3,1)_{-\frac{1}{2}}$	$(3,1)_{\frac{1}{2}}$	$(3,1)_{-\frac{3}{2}}$	$(3,2)_{\frac{1}{2}}$	$(3,2)_{\frac{3}{2}}$	$(3,3)_{-\frac{1}{2}}$

Name	Ω_1	Ω_2	Ω_4	Υ	Φ
Irrep	$(6,1)_{\frac{1}{2}}$	$(6,1)_{-\frac{1}{2}}$	$(6,1)_{\frac{3}{2}}$	$(6,3)_{\frac{1}{2}}$	$(8,2)_{\frac{1}{2}}$


Table 1. New scalar bosons contributing to the dimension-six SMEFT at tree level.

[LA, McCullough, Renner 2408.03992]

Fermions

(*) = special choice of couplings to avoid tree-level EWPO


■ Universal couplings ■ Third-gen. only ■ Other



(*) = special choice of couplings to avoid tree-level EWPO

■ Universal couplings ■ Third-gen. only ■ Flavourless couplings

▨ = no running



Flavour prospects at FCC-ee

Table 6: Yields of heavy-flavoured particles produced at FCC-ee for 6×10^{12} Z decays [190].

Particle species	B^0	B^+	B_s^0	Λ_b	B_c^+	$c\bar{c}$	$\tau^-\tau^+$
Yield ($\times 10^9$)	370	370	90	80	2	720	200

- > $\sim 10^3$ more $b\bar{b}$ and $\tau^+\tau^-$ w.r.to Belle
- > $\sim \times 5$ improvement in Λ_{NP} reach
- > ~ 30 more than Belle-II projections
- > Access to B_s and B_c - not produced at b factories
- > Great advantage due to clean environment and boosted final states

Attribute	$\Upsilon(4S)$	pp	Z
All hadron species		✓	✓
High boost		✓	✓
Enormous production cross-section		✓	(✓)
Negligible trigger losses	✓		✓
High geometrical acceptance	✓		✓
Low backgrounds	✓		✓
Flavour-tagging power	✓		✓
Initial-energy constraint	✓		(✓)

[Kamenik et al. '25]

Projections for flavour observables

[LA, Isidori, Pešut 2503.17019]

Observable	SM	Current value [14]	Pre-FCC projection	FCC-ee expected
$ g_\tau/g_\mu $	1	1.0009 ± 0.0014	–	± 0.0001 [15]
$ g_\tau/g_e $	1	1.0027 ± 0.0014	–	± 0.0001 [15]
corr.		0.51		
$\mathcal{B}(\tau \rightarrow \mu \bar{\mu} \mu)$	0	$< 2.1 \times 10^{-8}$	$< 0.37 \times 10^{-8}$ [*] [16]	$< 1.5 \times 10^{-11}$ [*] [15]
R_D	0.298 ± 0.004	0.342 ± 0.026 [17]	$\pm 3.0\%$ [16]	
R_{D^*}	0.254 ± 0.005	0.287 ± 0.012 [17]	$\pm 1.8\%$ [16]	
corr.		-0.39		
$\mathcal{B}(B_c \rightarrow \tau \bar{\nu})$	$(1.95 \pm 0.09) \times 10^{-2}$	< 0.3 (68% C.L.)	–	$\pm 1.6\%$ [8]
$\mathcal{B}(B \rightarrow K \nu \bar{\nu})$	$(4.44 \pm 0.30) \times 10^{-6}$	$(1.3 \pm 0.4) \times 10^{-5}$	$\pm 14\%$ [16]	$\pm 3\%$ [7]
$\mathcal{B}(B \rightarrow K^* \nu \bar{\nu})$	$(9.8 \pm 1.4) \times 10^{-6}$	$< 1.2 \times 10^{-5}$ (68% C.L.)	$\pm 33\%$ [16]	$\pm 3\%$ [7]
$\mathcal{B}(B \rightarrow K \tau \bar{\tau})$	$(1.42 \pm 0.14) \times 10^{-7}$	$< 1.5 \times 10^{-3}$ (68% C.L.)	$< 2.7 \times 10^{-4}$	$\pm 20\%$ [**] [18]
$\mathcal{B}(B \rightarrow K^* \tau \bar{\tau})$	$(1.64 \pm 0.06) \times 10^{-7}$	$< 2.1 \times 10^{-3}$ (68% C.L.)	$< 6.5 \times 10^{-4}$ [*] [16]	$\pm 20\%$ [**] [18]
$\mathcal{B}(B_s \rightarrow \tau \bar{\tau})$	$(7.45 \pm 0.26) \times 10^{-7}$	$< 3.4 \times 10^{-3}$ (68% C.L.)	$< 4.0 \times 10^{-4}$ [*] [16]	$\pm 10\%$ [**] [18]
$\Delta M_{B_s}/\Delta M_{B_s}^{\text{SM}}$	1	$\pm 7.6\%$	$\pm 3.3\%$ [19]	$\pm 1.5\%$ [19]
$\mathcal{B}(B \rightarrow K \tau \bar{\mu})$	0		$< 1.0 \times 10^{-6}$ [*] [20]	
$\mathcal{B}(B_s \rightarrow \tau \bar{\mu})$	0		$< 1.0 \times 10^{-6}$ [*] [20]	

> Subset of observables, relevant for our example study

[**] = under the assumption of an enhanced rate due to NP

Third-gen. semileptonics: future prospects


SMEFT analysis

- > Current tension driven by $R_{D^{(*)}}$
- > Only third-gen. flavour indices:
$$\mathcal{L} \supset -\frac{2}{v^2} (\bar{\ell}_L^3 \sigma^I \gamma_\mu \ell_L^3) (\bar{q}_L^3 \sigma^I \gamma^\mu q_L^3)$$
- > Flavour-violating effects via $U(2)_q$ breaking spurion

$$\tilde{V} = -\varepsilon V_{ts} \begin{pmatrix} V_{td}/V_{ts} \\ 1 \end{pmatrix} \quad \varepsilon \sim \mathcal{O}(1)$$

- > $q_L^3 \rightarrow q_L^3 + \tilde{V}_i q_L^i$
- > Assume a signal compatible with current measurements (grey region), and project for FCC-ee expected errors

[LA, Isidori, Pešut 2503.17019]



Flavour v. EWPOs


[LA, Isidori, Pešut 2503.17019]

$$\mathcal{O}_{\ell q}^{(1)} = (\bar{\ell}_L^3 \gamma_\mu \ell_L^3)(\bar{q}_L^3 \gamma^\mu q_L^3)$$

$$\mathcal{O}_{\ell q}^{(3)} = (\bar{\ell}_L^3 \sigma^I \gamma_\mu \ell_L^3)(\bar{q}_L^3 \sigma^I \gamma^\mu q_L^3)$$

- > Flavour-conserving directions (no spurions) badly probed by flavour
- > Interplay with EWPOs: independent probes
- > y_t running into $C_{H\ell}^{(1\pm 3)}$

Complementarity!




An explicit (simplified) model: S_1 scalar leptoquark



Synergies between Higgs/EW/Flavour

$$S_1 \sim (\mathbf{3}, \mathbf{1}, 1/3)$$

$$\mathcal{L}_{S_1} \supset i y_L S_1 (\bar{q}_L^c \sigma_2 \ell_L^3) + y_R S_1 (\bar{u}^3 e_R^3) + \text{h.c.}$$



- > Third-generation couplings only
- > Current data



$$q_L^3 \rightarrow q_L^3 - \epsilon_S V_{ti} q_L^i$$

An explicit (simplified) model: S_1 scalar leptoquark



Synergies between Higgs/EW/Flavour

$$S_1 \sim (\mathbf{3}, \mathbf{1}, 1/3)$$

$$\mathcal{L}_{S_1} \supset i y_L S_1 (\bar{q}_L^c \sigma_2 \ell_L^3) + y_R S_1 (\bar{u}^3 e_R^3) + \text{h.c.}$$



- > Third-generation couplings only
- > Current data
- > $H \rightarrow \tau\tau$ (1%)




$$q_L^3 \rightarrow q_L^3 - \epsilon_S V_{ti} q_L^i$$

An explicit (simplified) model: S_1 scalar leptoquark



Synergies between Higgs/EW/Flavour

$$S_1 \sim (\mathbf{3}, \mathbf{1}, 1/3)$$

$$\mathcal{L}_{S_1} \supset i y_L S_1 (\bar{q}_L^c \sigma_2 \ell_L^3) + y_R S_1 (\bar{u}^3 e_R^3) + \text{h.c.}$$



- > Third-generation couplings only
- > Current data
- > $H \rightarrow \tau\tau$ (1%)
- > A_τ ($\sim 10^{-4}$)




$$q_L^3 \rightarrow q_L^3 - \epsilon_S V_{ti} q_L^i$$

An explicit (simplified) model: S_1 scalar leptoquark



Synergies between Higgs/EW/Flavour

$$\mathcal{L}_{S_1} \supset i y_L S_1 (\bar{q}_L^c \sigma_2 \ell_L^3) + y_R S_1 (\bar{u}^3 e_R^3) + \text{h.c.} \quad S_1 \sim (\mathbf{3}, \mathbf{1}, 1/3)$$

[LA, Isidori, Pešut 2503.17019]



- > Third-generation couplings only
- > Current data
- > $H \rightarrow \tau\tau$ (1%)
- > A_τ ($\sim 10^{-4}$)
- > $B \rightarrow K\nu\bar{\nu}$ (3%)



$$q_L^3 \rightarrow q_L^3 - \epsilon_S V_{ti} q_L^i$$

EWPOs off the pole: four-fermion interactions

Energy-enhancement compensates luminosity

- > Four-fermion operators, e.g. $e\bar{e}bb$, have negligible contribution on the pole to e.g. R_b
- > Off the pole, energy-enhanced amplitudes in the EFT


$$\mathcal{A}(e^+e^- \rightarrow f\bar{f}) \sim \frac{s}{\Lambda^2} \mathcal{C}_{eeff}$$

- > Comparable sensitivity w.r.to the pole (loop effects)
- > Can dominate when the RGE effect is smaller (e.g. gauge coupling running)
- > Define ratios

$$R_a = \frac{\sigma(e^+e^- \rightarrow a\bar{a})}{\sum_{q=u,d,s,c,b} \sigma(e^+e^- \rightarrow q\bar{q})}$$

- > At $t\bar{t}$, define $R_t = \sigma(e^+e^- \rightarrow t\bar{t})/\sigma(e^+e^- \rightarrow q\bar{q})$


[Greljo, Tiplom, Valenti 2411.02485]



EWPOs off the pole: four-fermion interactions

Energy-enhancement compensates luminosity



[Greljo, Tiblom, Valenti 2411.02485]



The Higgs trilinear at FCC-ee

Global fits with flavour symmetries

- > Measure κ_λ from $e^+e^- \rightarrow Zh$
- > Expected precision $\sim 20\%$




[Maura, Stefaneke, You 2503.13719]


- > See also ter Hoeve et al. (SMEFiT) 2504.05974

Trilinear v. electron-top interactions

- > C_H and $eett$ interactions enter at the same order in $e^+e^- \rightarrow Zh$
- > Possibly spoil the measurement of λ_3
- > Can we independently constrain $eett$ operators? What's the impact on the trilinear?



$$\mathcal{O}_H = (H^\dagger H)^3$$




$$[\mathcal{O}_{lq}^{(1)}]_{1133} = (\bar{l}_1 \gamma_\mu l_1)(\bar{q}_3 \gamma^\mu q_3),$$

$$[\mathcal{O}_{lq}^{(3)}]_{1133} = (\bar{l}_1 \gamma_\mu \sigma^I l_1)(\bar{q}_3 \gamma^\mu \sigma^I q_3),$$

$$[\mathcal{O}_{qe}]_{3311} = (\bar{q}_3 \gamma^\mu q_3)(\bar{e}_1 \gamma_\mu e_1),$$

Trilinear v. electron-top interactions

[LA, Grojean, Tabatt 2512.06916]










Higgs on the Z -pole

[Maura, Stefaneke, You 2412.14241]

Precision against loops

> EWPOs on the Z -pole can be sensitive and complementary probes





Measuring the electron Yukawa?

EFT perspective

- > At a dedicated Higgs-pole run (~ 1 yr), get $\kappa_e \lesssim 1.6$
- > In SMEFT, $\mathcal{L} \supset \mathcal{C}_{eH}(\bar{\ell}eH)(H^\dagger H)$

$$\kappa_e = 1 - \frac{v^3}{\sqrt{2} m_e} \mathcal{C}_{eH}.$$

- > Two-loop connection with the electron
 $g - 2$ unavoidable





[LA, McCullough, Renner, Rocha, Smith 2511.02642]

Measuring the electron Yukawa?

Models

- > In explicit models, relation between $(g - 2)_e$ and y_e can change
- > e.g. with vector-like leptons



[Erdelyi, Gröber, Selimović 2501.07628]

Summary

- > Indirect searches for New Physics can probe much higher scales than direct searches
- > FCC-ee, with a leap in precision, will provide an unprecedented opportunity for precision physics
- > EWPOs can probe many different NP scenarios
→ sensitive to loop effects from heavy NP
- > Linear extensions of the SM probed from $\mathcal{O}(1)$ up to $\mathcal{O}(100)$ TeV scale
- > (Heavy) flavour prospects complementary to B factories and LHC
- > Higgs physics probed from different angles (here: κ_λ , κ_e)
- > Great potential for TeV-scale, third-gen. NP
- > Exciting precision program overall - still much to be explored!

Thank you!


lukas.allwicher@desy.de

Backup



Probing the Granada dictionary at FCC-ee

SMEFiT collaboration 2502.20453



Higgs prospects at FCC-ee

FCC FSR 2505.00272

Coupling	HL-LHC	FCC-ee	FCC-ee + FCC-hh
κ_Z (%)	1.3*	0.10	0.10
κ_W (%)	1.5*	0.29	0.25
κ_b (%)	2.5*	0.38 / 0.49	0.33 / 0.45
κ_g (%)	2*	0.49 / 0.54	0.41 / 0.44
κ_τ (%)	1.6*	0.46	0.40
κ_c (%)	—	0.70 / 0.87	0.68 / 0.85
κ_γ (%)	1.6*	1.1	0.30
$\kappa_{Z\gamma}$ (%)	10*	4.3	0.67
κ_t (%)	3.2*	3.1	0.75
κ_μ (%)	4.4*	3.3	0.42
$ \kappa_S $ (%)	—	$^{+29}_{-67}$	$^{+29}_{-67}$
Γ_H (%)	—	0.78	0.69
$\mathcal{B}_{\text{inv}} (<, 95\% \text{ CL})$	$1.9 \times 10^{-2} *$	5×10^{-4}	2.3×10^{-4}
$\mathcal{B}_{\text{unt}} (<, 95\% \text{ CL})$	$4 \times 10^{-2} *$	6.8×10^{-3}	6.7×10^{-3}

The Standard Model flavour puzzle

- SM gauge interactions are flavour-blind:

$$\mathcal{L}_{\text{gauge}} \supset \sum_{i=1}^3 \sum_{\psi} \bar{\psi}_i i \not{D} \psi_i$$

- Flavour symmetry:

$$U(3)^5 = U(3)_q \times U(3)_u \times U(3)_d \times U(3)_\ell \times U(3)_e$$

- Only breaking: Higgs Yukawa interactions:

$$\mathcal{L}_{\text{Yukawa}} = Y_u \bar{q} u \tilde{H} + Y_d \bar{q} d H + Y_e \bar{\ell} e H + \text{h.c.}$$



The pattern of masses and mixings doesn't look accidental at all!

$$m_{\psi} \sim \begin{pmatrix} \text{light} & & \\ & \text{medium} & \\ & & \text{heavy} \end{pmatrix} \quad V_{\text{CKM}} \sim \begin{pmatrix} \text{heavy} & \text{medium} & \\ \text{medium} & \text{heavy} & \text{light} \\ & \text{light} & \text{heavy} \end{pmatrix}$$

Example: $d_i \rightarrow d_j \nu \nu$ transitions

- > FCNC: high-scale probes due to loop+GIM suppression
- > Clean: no theory uncertainty due to charm rescattering
- > Very rare, exp. challenging
- > Only probes of third-gen. leptons so far
- > First measurement in 2023 by Belle-II:

> NA62 2024



SMEFT description of $d_i \rightarrow d_j \nu \bar{\nu}$

[1903.10954]

- > Start with third-generation indices only: **rank-one hypothesis**

$$Q_{\ell q}^{\pm} = (\bar{q}_L^3 \gamma^{\mu} q_L^3)(\bar{\ell}_L^3 \gamma_{\mu} \ell_L^3) \pm (\bar{q}_L^3 \gamma^{\mu} \sigma^a q_L^3)(\bar{\ell}_L^3 \gamma_{\mu} \sigma^a \ell_L^3)$$

$$Q_S = (\bar{\ell}_L^3 \tau_R)(\bar{b}_R q_L^3)$$

- > Third generation LH quarks: **down alignment**


$$q_L^3 = \begin{pmatrix} V_{ub}u_L + V_{cb}c_L + V_{tb}t_L \\ b_L \end{pmatrix}$$

- > $U(2)_q$ -breaking spurion

$$\tilde{V} = -\varepsilon V_{ts} \begin{pmatrix} \kappa V_{td}/V_{ts} \\ 1 \end{pmatrix}$$

- > Replace $q_L^3 \rightarrow q_L^3 + \tilde{V}_i q_L^i$
- > System described by 5 parameters: $C_S, C_{\ell q}^+, C_{\ell q}^-, \varepsilon, \kappa$

Correlated observables




	C_S	$C_{\ell q}^+$	$C_{\ell q}^-$	ε	κ	Exp. indication
$\sigma(pp \rightarrow \ell\ell)$	✓	✓	✓			bounds on $\mathcal{A}_{\text{NP}}$
EWPO		✓	✓			bounds on $\mathcal{A}_{\text{NP}}$
R_D, R_{D^*}	✓	✓	✓	✓		$\mathcal{A}_{\text{NP}}/\mathcal{A}_{\text{SM}} > 0$
$\mathcal{B}(B \rightarrow K^{(*)} \mu \bar{\mu})$		✓		✓		$\mathcal{A}_{\text{NP}}/\mathcal{A}_{\text{SM}} < 0$
$\mathcal{B}(B \rightarrow K \nu \bar{\nu})$			✓	✓		$ \mathcal{A}_{\text{SM}} + \mathcal{A}_{\text{NP}} ^2 > \mathcal{A}_{\text{SM}} ^2$
$\mathcal{B}(K \rightarrow \pi \nu \bar{\nu})$			✓	✓	✓	$ \mathcal{A}_{\text{SM}} + \mathcal{A}_{\text{NP}} ^2 > \mathcal{A}_{\text{SM}} ^2$

Results: $C_{\ell q}^+ - \varepsilon$

- > Global fit without di-neutrino modes (don't affect $C_{\ell q}^+$)
- > LHC Drell-Yan + EWPO provide constraints on $C_{\ell q}^\pm$ and C_S
- > C_S compatible with zero (LHC constraints strong)
- > Non-zero $C_{\ell q}^+$ and ε driven by $R_{D^{(*)}}$:

$$\begin{aligned} \frac{R_{D^{(*)}}}{R_{D^{(*)}}^{\text{SM}}} &\approx 1 + 2\text{Re}(C_{V_L}) \\ &\approx 1 - v^2 (1 + \varepsilon) (C_{\ell q}^+ - C_{\ell q}^-) \end{aligned}$$

- > Suppress $|\varepsilon| > 3$ with theoretical likelihood




Results: $C_{\ell q}^- - \varepsilon$

- > Grey: Global fit without di-neutrino modes, $\kappa = 1$
- > $C_{\ell q}^-$ largely unconstrained
- > Good compatibility with di-neutrino modes for $\kappa = 1$

$$|C_{\tau,bs}^{\text{SM}}| \rightarrow \left| C_{\tau,bs}^{\text{SM}} - \varepsilon \frac{\pi v^2}{\alpha} C_{\ell q}^- \right|,$$


$$|C_{\tau,sd}^{\text{SM}}| \rightarrow \left| C_{\tau,sd}^{\text{SM}} + \kappa \varepsilon^2 \frac{\pi v^2}{\alpha} C_{\ell q}^- \right|.$$

- > Select $C_{\ell q}^- > 0$



Multi-scale picture

[Dvali, Shifman '00, 1603.06609, ..., 1712.01368, 2103.15635, 2303.01520, 2305.16280]



- **Idea:** non-universal interactions, with different scales coupling to different fermion families
- $\Lambda_1 \gg \Lambda_2 \gg \Lambda_3 \gg m_W$
- Gauge universality appears as an accident at low energies
- Good way of explaining fermion hierarchies
- Don't decouple flavour-universal NP from a high-scale explanation of the fermion hierarchies
- $\Lambda_3 \sim \text{TeV}$ helps with the Higgs hierarchy problem

Phenomenology: Flavour

- In principle, no flavour-violating couplings in the exact $U(2)$ limit
- But, need to specify a basis for the quark doublets
- Two choices:

down-aligned

$$q_L^{\text{down}} = \begin{pmatrix} V_{\text{CKM}}^\dagger u_L \\ d_L \end{pmatrix}$$

up-aligned

$$q_L^{\text{up}} = \begin{pmatrix} u_L \\ V_{\text{CKM}} d_L \end{pmatrix}$$

- Main effects in the $U(2)$ hypothesis:
 - $\Delta F = 1$: $B \rightarrow X_s \gamma$, $B \rightarrow K \nu \bar{\nu}$, $K \rightarrow \pi \nu \bar{\nu}$, $B \rightarrow K \mu \mu$, ...
 - $\Delta F = 2$: B_{s,d^-} , K^- , D -mixing
 - $b \rightarrow c \tau \nu$ transitions: $R_{D^{(*)}}$, R_{Λ_c}


Phenomenology: EWPO

- Crucial precision tests of the SM & NP coupled to the Higgs
- At tree-level, constrain operators of the type $(H^\dagger i D_\mu H)(\bar{\psi} \gamma^\mu \psi)$
 - modification of SM gauge boson couplings
 - Only 15 such structures in the $U(2)^5$ limit
- Include also Higgs decays, τ LFU tests

#	Wilson Coef.	[Obs] bound	Λ_{bound} [TeV]
1	cHWB	A_b^{FB}	9.63
2	CHl1[l]	σ_{had}	8.07
3	CHl3[l]	A_b^{FB}	7.96
4	CHe[l]	σ_{had}	6.93
5	cHD	A_b^{FB}	5.74
6	CHq3[l]	R_c	5.73
7	CHl1[h]	R_c	4.57
8	CHl3[h]	R_c	4.48
9	Cl1[l, p, p, l]	A_b^{FB}	4.43
10	CHe[h]	R_c	3.97
11	CHq3[h]	R_b	3.43
12	CHq1[h]	R_b	3.43
13	CHu[l]	R_c	2.58
14	CHq1[l]	R_c	2.07
15	CHd[l]	R_c	1.81
16	CHd[h]	R_b	1.4

Phenomenology: colliders

- High- p_T Drell-Yan Tails



- In particular: $pp \rightarrow \tau\tau, \tau\nu$
- Constrain semileptonic operators

[LA, Faroughy, Jaffredo, Sumensari, Wilsch 2207.10756]

- LEP-2 $e^+e^- \rightarrow \ell^+\ell^-$

- $e^+e^- \rightarrow e^+e^-$ angular distributions
- $e^+e^- \rightarrow \mu^+\mu^-, \tau^+\tau^-$: $\sigma, \sigma_{\text{FB}}$
- Constrain four-lepton operators

[Allanach, Mullin 2306.80669]

- four-quark observables

- $t\bar{t}, b\bar{b}, b\bar{t}$ final states
- Constrain e.g. $\mathcal{C}_{qu}^{(1)}, \mathcal{C}_{uG}, \dots$

[Ethier et al. 2105.00006]



Suppressing the light families

[LA, Cornella, Isidori, Stefaneke 2311.00020]

- So far, only $U(2)^5$ protection
- No suppression of operators involving the light families
- ε_Q for each light quark field
- ε_L for each light lepton field


Examples:

$$C_{He}^{[ii]}(H^\dagger i \overleftrightarrow{D}_\mu H) \sum_{i=1}^2 (\bar{e}_i \gamma^\mu e_i) \rightarrow \varepsilon_L^2 C_{He}^{[ii]}(H^\dagger i \overleftrightarrow{D}_\mu H) \sum_{i=1}^2 (\bar{e}_i \gamma^\mu e_i)$$
$$C_{\ell q}^{(1)[ijjj]} \sum_{i,j=1}^2 (\bar{\ell}^i \gamma^\mu \ell^i)(\bar{q}^j \gamma_\mu q^j) \rightarrow \varepsilon_L^2 \varepsilon_Q^2 C_{\ell q}^{(1)[ijjj]} \sum_{i,j=1}^2 (\bar{\ell}^i \gamma^\mu \ell^i)(\bar{q}^j \gamma_\mu q^j)$$

- Dial down ε_i until collider bounds are below $\Lambda_0 = 1.5$ TeV

Suppressing the light families


[LA, Cornella, Isidori, Stefaneke 2311.00020]




- operators with Higgs fields still give strong bounds (EWPO)

The Higgs and $U(2)^5$

- If we want to address both the Higgs hierarchy problem and the flavour puzzle, NP should couple to the Higgs as well
- Take e.g. a Z' model, one generically gets contributions to EWPO



$$C_{Hq}^{(1)[33]} (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_L^3 \gamma^\mu q_L^3)$$




$$C_{HD} |H^\dagger \overleftrightarrow{D}_\mu H|^2$$

$U(2)^5$ does not offer protection for these contributions

- Need to suppress the NP couplings to the Higgs to avoid EWPO constraints
- ε_H for each Higgs field in the EFT

Suppressing Higgs couplings

[LA, Cornella, Isidori, Stefaneke 2311.00020]



- Some flavour bounds still large (in the up-aligned case)

Flavour alignment in the 3rd generation

[LA, Cornella, Isidori, Stefaneke 2311.00020]

- q_L^3 is somewhere in-between down-aligned and up-aligned
- ε_F to parametrise the amount of down-alignment:

$$\theta \sim V_{cb}\varepsilon_F$$

$$\begin{pmatrix} t_L \\ V_{td}d_L + V_{ts}s_L + V_{tb}b_L \end{pmatrix} = q_t \uparrow \quad \begin{array}{c} q_3 \\ \nearrow V_{cb} \\ \nearrow \varepsilon_F V_{cb} \\ \nearrow q_b \end{array} \quad q_b = \begin{pmatrix} V_{ub}^*u_L + V_{cb}^*c_L + V_{tb}^*t_L \\ b_L \end{pmatrix}$$

$$\begin{aligned} q_3 &= [(1 - \varepsilon_F)\delta_{3r} + \varepsilon_F V_{3r}]q_r^{(d)} \approx q_b + \varepsilon_F(V_{ts}q_s + V_{td}q_d) \\ &= [(1 - \varepsilon_F)(V^\dagger)_{3r} + \varepsilon_F\delta_{3r}]q_r^{(u)} \approx \varepsilon_F q_t + (1 - \varepsilon_F)(V_{cb}^*q_c + V_{ub}^*q_u) \end{aligned}$$

SM predictions: impact of $|V_{cb}|$

$$O_{\ell,ij}^\nu = (\bar{d}_L^i \gamma_\mu d_L^j)(\bar{\nu}_L^\ell \gamma^\mu \nu_L^\ell)$$

- > At $\mu = m_{b,s}$: $\mathcal{L}_{\text{eff}} = \frac{4G_F}{\sqrt{2}} \frac{\alpha}{2\pi} \sum_{\ell=e,\mu,\tau} \left[\lambda_{sd}^t C_{\ell,sd}^{\text{SM}} O_{\ell,sd}^\nu + \lambda_{bs}^t C_{\ell,bs}^{\text{SM}} O_{\ell,bs}^\nu \right] + \text{h.c.}$
- > Leading uncertainty from V_{cb} :

$$\lambda_{sd}^t = V_{ts} V_{td}^* = \lambda |V_{cb}|^2 \left[(\bar{\rho} - 1) \left(1 - \frac{\lambda^2}{2} \right) + i\bar{\eta} \left(1 + \frac{\lambda^2}{2} \right) \right]$$

- > Take average between inclusive and exclusive, inflating errors [Finauri+Gambino '24]
[Bordone+Juttner '24]


- > First measurement by NA62 in 2024!

$$|V_{cb}|_{\text{incl+excl}} = (41.37 \pm 0.81) \times 10^{-3}$$

$$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})^{\text{SM}} = (8.09 \pm 0.63) \times 10^{-11}$$


- > $b \rightarrow s \nu \bar{\nu}$: Bečirević et al. 2301.06990

$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu}) / |\lambda_{bs}^t|^2 = (2.87 \pm 0.10) \times 10^{-3}$$




Future prospects

> Measure ε with dineutrino modes



> Minimal vs. non-minimal $U(2)_q$ breaking



Z- and W Lagrangian

- Effective description at the EW scale:

$$\begin{aligned} \mathcal{L}_{\text{eff}} \supset & -\frac{g_L}{\sqrt{2}} W^{+\mu} \left[\bar{u}_L^i \gamma_\mu \left(V_{ij} + \delta g_{ij}^{Wq} \right) d_L^j + \bar{\nu}_L^i \gamma_\mu \left(\delta_{ij} + \delta g_{ij}^{W\ell} \right) e_L^j \right] + \text{h.c.} \\ & -\sqrt{g_L^2 + g_Y^2} Z^\mu \left(\bar{f}_L^i \gamma_\mu \left(g_L^{Zf} \delta_{ij} + \delta g_{Lij}^{Zf} \right) f_L^j + \bar{f}_R^i \gamma_\mu \left[g_R^{Zf} \delta_{ij} + \delta g_{Rij}^{Zf} \right] f_R^j \right) \\ & + \frac{g_L^2 v^2}{4} (1 + \delta m_W)^2 W^{+\mu} W_\mu^- + \frac{g_L^2 v^2}{8c_W^2} Z^\mu Z_\mu \end{aligned}$$

- Given already current exp. precision, expect small NP effects
→ consider only flavour-conserving pieces (interference with SM)
- $SU(2)_L$ fixes

$$\delta g_L^{Z\nu} = \delta g_L^{W\ell} + \delta g_L^{Ze}, \quad \delta g_L^{Wq} = \delta g_L^{Zu} V - V \delta g_L^{Zd}$$

- 20 independent parameters:

$$\delta g \in \{ \delta g_{L11(22,33)}^{Z\nu}, \delta g_{L11(22,33)}^{Ze}, \delta g_{R11(22,33)}^{Ze}, \delta g_{L11(22)}^{Zu}, \delta g_{R11(22)}^{Zu}, \delta g_{L11(22,33)}^{Zd}, \delta g_{R11(22,33)}^{Zd}, \delta m_W \}$$

Mapping to SMEFT: basis

- Use Warsaw basis [1008.4884]
- $\psi^2 H^2 D$ operators

$$\mathcal{C}_{H\psi}(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{\psi}\gamma^\mu\psi) \xrightarrow{H \rightarrow \langle H \rangle} -\frac{v^2}{2}\mathcal{C}_{H\psi}Z_\mu(\bar{\psi}\gamma^\mu\psi)$$

$[\mathcal{O}_{H\ell}^{(1)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{\ell}_p \gamma^\mu \ell_r)$	$[\mathcal{O}_{He}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$[\mathcal{O}_{H\ell}^{(3)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{\ell}_p \tau^I \gamma^\mu \ell_r)$	$[\mathcal{O}_{Hu}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
$[\mathcal{O}_{Hq}^{(1)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$	$[\mathcal{O}_{Hd}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$[\mathcal{O}_{Hq}^{(3)}]_{pr} = (H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$	

- Bosonic operators

$$\mathcal{O}_{HD} = |H^\dagger D_\mu H|^2 \quad \mathcal{O}_{HWB} = (H^\dagger \tau^I H) W_{\mu\nu}^I B^{\mu\nu}$$

→ associated with T and S parameters

- Four-fermion operator

$$[\mathcal{O}_{\ell\ell}]_{1221} = (\bar{\ell}_1 \gamma^\mu \ell_2)(\bar{\ell}_2 \gamma^\mu \ell_1)$$

Mapping to SMEFT: matching

- Matching of the δg s:

$$\delta g_{Lii}^{Z\nu} = -\frac{v^2}{2} \left([\mathcal{C}_{H\ell}^{(1)}]_{ii} - [\mathcal{C}_{H\ell}^{(3)}]_{ii} \right) + \delta^U(1/2, 0)$$

$$\delta g_{Lii}^{Ze} = -\frac{v^2}{2} \left([\mathcal{C}_{H\ell}^{(1)}]_{ii} + [\mathcal{C}_{H\ell}^{(3)}]_{ii} \right) + \delta^U(-1/2, -1)$$

$$\delta g_{Rii}^{Ze} = -\frac{v^2}{2} [\mathcal{C}_{He}]_{ii} + \delta^U(0, -1)$$

$$\delta g_{Lii}^{Zu} = -\frac{v^2}{2} \left([\mathcal{C}_{Hq}^{(1)}]_{ii} - [\mathcal{C}_{Hq}^{(3)}]_{ii} \right) + \delta^U(1/2, 2/3),$$

$$\delta g_{Rii}^{Zu} = -\frac{v^2}{2} [\mathcal{C}_{Hu}]_{ii} + \delta^U(0, 2/3)$$

$$\delta g_{Lii}^{Zd} = -\frac{v^2}{2} \left([\mathcal{C}_{Hq}^{(1)}]_{ii} + [\mathcal{C}_{Hq}^{(3)}]_{ii} \right) + \delta^U(-1/2, -1/3)$$

$$\delta g_{Rii}^{Zd} = -\frac{v^2}{2} [\mathcal{C}_{Hd}]_{ii} + \delta^U(0, -1/3)$$

$$\delta m_W = -\frac{v^2 g_L^2}{4(g_L^2 - g_Y^2)} \mathcal{C}_{HD} - \frac{v^2 g_L g_Y}{g_L^2 - g_Y^2} \mathcal{C}_{HWB} + \frac{v^2 g_Y^2}{4(g_L^2 - g_Y^2)} \left([\mathcal{C}_{\ell\ell}]_{1221} - 2[\mathcal{C}_{H\ell}^{(3)}]_{22} - 2[\mathcal{C}_{H\ell}^{(3)}]_{11} \right)$$

- Universal contribution:

$$\delta^U(T^3, Q) = -v^2 \left(T^3 + Q \frac{g_Y^2}{g_L^2 - g_Y^2} \right) \left(\frac{1}{4} \mathcal{C}_{HD} + \frac{1}{2} [\mathcal{C}_{H\ell}^{(3)}]_{22} + \frac{1}{2} [\mathcal{C}_{H\ell}^{(3)}]_{11} - \frac{1}{4} [\mathcal{C}_{\ell\ell}]_{1221} \right) - v^2 Q \frac{g_L g_Y}{g_L^2 - g_Y^2} \mathcal{C}_{HWB}$$

Input redefinitions

- Universal contribution:

$$\delta^U(T^3, Q) = -v^2 \left(T^3 + Q \frac{g_Y^2}{g_L^2 - g_Y^2} \right) \left(\frac{1}{4} \mathcal{C}_{HD} + \frac{1}{2} [\mathcal{C}_{H\ell}^{(3)}]_{22} + \frac{1}{2} [\mathcal{C}_{H\ell}^{(3)}]_{11} - \frac{1}{4} [\mathcal{C}_{\ell\ell}]_{1221} \right) - v^2 Q \frac{g_L g_Y}{g_L^2 - g_Y^2} \mathcal{C}_{HWB}$$

- Input scheme: α, G_F, m_Z
- G_F is extracted from $\mu \rightarrow e \nu \bar{\nu}$ decay
→ anything modifying the decay rate in SMEFT is not observable, but gets propagated through the inputs (vev, gauge couplings)
 - $[\mathcal{C}_{H\ell}^{(3)}]_{11,22}$: W -coupling modifications
 - $[\mathcal{C}_{\ell\ell}]_{1221}$: direct four-fermion contact interaction
- m_Z gets a contribution from $\mathcal{C}_{HD}$ and $\mathcal{C}_{HWB}$ in SMEFT

The set of Electroweak Wilson Coefficients

- From matching conditions + input redefinitions:

$$\vec{C}_{\text{ew}} = ([C_{H\ell}^{(1,3)}]_{ii}, [C_{Hq}^{(1,3)}]_{ii}, [C_{He}]_{ii}, [C_{Hu}]_{ii}, [C_{Hd}]_{ii}, C_{HD}, C_{HWB}, [C_{\ell\ell}]_{1221})$$

- Note: For C_{Hu} $i = 1, 2$ only (no RH tops)
- Total of **23 parameters**
- Assuming all 20 δg s can be constrained individually, one expects 3 flat directions in the SMEFT EW fit

$$C_0^{(1)} \propto [C_{Hq}^{(1)}]_{33} - [C_{Hq}^{(3)}]_{33}$$

$$C_0^{(2)} \propto -\frac{g_Y}{g_L} C_{HWB} + \sum_{i=1}^3 \left([C_{H\ell}^{(3)}]_{ii} + [C_{Hq}^{(3)}]_{ii} \right) \quad [\text{cf. also } \text{hep-ph}/0602154 \text{ and } 1701.06424]$$

$$C_0^{(3)} \propto 2C_{HD} - \frac{1}{2} \frac{g_L}{g_Y} C_{HWB} + \sum_{\psi} \sum_i Y_{\psi} [C_{H\psi}]_{ii}$$

RGE effects, systematically

- RGE equations for EW operators (y_t , g_1 and g_2 contributions)
- Get sensitivity to 168 operators

$$\begin{aligned}
 \dot{C}_{HD} &= N_c \left(\frac{16}{3} Y_H Y_u g_1^2 - 8 y_t^2 \right) C_{Hu}^{33} + \frac{80}{3} g_1^2 Y_H^2 C_{H\Box} \\
 \dot{C}_{Hf_1}^{(1)jj} &= 2 N_c y_t^2 \left(-S_{f_1 u} C_{uf_1}^{33jj} + S_{f_1 q} C_{qf_1}^{33jj} \right) \\
 \dot{C}_{Hl}^{(3)jj} &= -2 N_c y_t^2 C_{lq}^{(3)jj33} \\
 \dot{C}_{Hq}^{(3)jj} &= -2 y_t^2 \left(2 N_c C_{qq}^{(3)jj33} + C_{qq}^{(1)j33j} - C_{qq}^{(3)j33j} \right) \\
 \dot{C}_{Hf_1}^{(1)jj} &= \frac{4}{3} g_1^2 Y_h \sum_k \left[Y_e S_{f_1 e} C_{f_1 e}^{jjkk} + 2 Y_l S_{f_1 l} C_{f_1 l}^{jjkk} \right. \\
 &\quad \left. + N_c \left(Y_u S_{f_1 u} C_{f_1 u}^{jjkk} + Y_d S_{f_1 d} C_{f_1 d}^{jjkk} + 2 Y_q S_{f_1 q} C_{f_1 q}^{jjkk} \right) \right] \\
 \dot{C}_{Hl}^{(3)jj} &= \frac{2}{3} g_2^2 \sum_k \left(C_{ll}^{jjkk} + N_c C_{lq}^{(3)jjkk} \right), \\
 \dot{C}_{Hq}^{(3)jj} &= \frac{2}{3} g_2^2 \sum_k \left(C_{lq}^{(3)kkjj} + 2 N_c C_{qq}^{(3)kkjj} + C_{qq}^{(1)jkkj} - C_{qq}^{(3)jkkj} \right) \\
 \dot{C}_{ll}^{1221} &= \frac{2}{3} g_2^2 \left[C_{ll}^{2222} + C_{ll}^{2332} + C_{ll}^{1111} + C_{ll}^{1331} + N_c \sum_k \left(C_{lq}^{(3)22kk} + C_{lq}^{(3)11kk} \right) \right]
 \end{aligned}$$

To avoid EWPOs at one loop, need to have R.H.S. = 0 for all the above *at the same time*

Scalars

State	Tree level zero possible?	1-loop RGE zero possible?	Bounding obs.
$\mathcal{S}$	✓	✗	A_e
$\mathcal{S}_1$	✓ if $(y_{\mathcal{S}_1})_{12} = 0$	✗	A_e
$\mathcal{S}_2$	✓	✗	$A_e (R_\tau)$
φ	✓	✗	Γ_Z
Ξ	✗	✗	m_W
Ξ_1	✓ if $\kappa_{\Xi_1} = 0$ & $(y_{\Xi_1})_{12} = 0$	✗	m_W
Θ_1	✓	✓*	m_W
Θ_3	✓	✓*	m_W
ω_1	✓	✗	Γ_Z
ω_2	✓	✗	Γ_Z
ω_4	✓	✗	$A_e (R_\tau)$
Π_1	✓	✗	$A_e (R_\tau)$
Π_7	✓	✗	R_τ
ζ	✓	✗	$\Gamma_Z (R_\tau)$
Ω_1	✓	✗	Γ_Z
Ω_2	✓	✗	Γ_Z
Ω_4	✓	✓ if $(y_{\Omega_4})_{33} \neq 0$, all else zero	$\Gamma_Z (-)$
Υ	✓	✗	Γ_Z
Φ	✓	✗	Γ_Z

Fermions

State	Tree level zero possible?	1-loop RGE zero possible?	Bounding obs.
N	$\times$	$\times$	$A_e(\Gamma_Z)$
E	$\times$	$\times$	$\Gamma_Z(R_\tau)$
Δ_1	$\times$	$\times$	$A_e(R_\tau)$
Δ_3	$\times$	$\times$	$A_e(R_\tau)$
Σ	$\times$	$\times$	$\Gamma_Z(R_\tau)$
Σ_1	$\times$	$\times$	$A_e(\Gamma_Z)$
U	✓ if $(\lambda_U)_3 \neq 0$, all else zero	$\times$	$\Gamma_Z(m_W)$
D	$\times$	$\times$	Γ_Z
Q_1	✓ if $(\lambda_{Q_1}^u)_3 \neq 0$, all else zero	$\times$	m_W
Q_5	$\times$	$\times$	Γ_Z
Q_7	✓ if $(\lambda_{Q_7})_3 \neq 0$, all else zero	$\times$	m_W
T_1	$\times$	$\times$	$m_W(\Gamma_Z)$
T_2	$\times$	$\times$	Γ_Z



Vectors

State	Tree level zero possible?	1-loop RGE zero possible?	Bounding obs.
$\mathcal{B}$	✓ if $(g_B^\phi) = 0$ & $(g_B^l)_{12} = 0$	✓ if eqns. (??)	m_W
$\mathcal{B}_1$	✓ if $(g_{B_1}^\phi) = 0$	✗	m_W
$\mathcal{W}$	✓ if $(g_W^\phi) = 0$ & $K_{12}(g_W^l) = 0$	✓ if eqns. (??)	$A_e(\Gamma_Z)$
$\mathcal{W}_1$	✗	✗	m_W
$\mathcal{G}$	✓	✓ if $(g_G^u)_{33} \neq 0$, all else zero	Γ_Z
$\mathcal{G}_1$	✓	✗	Γ_Z
$\mathcal{H}$	✓	✗	Γ_Z
$\mathcal{L}_1$	✗	✗	A_e
$\mathcal{L}_3$	✓	✗	$A_e(A_\tau)$
$\mathcal{U}_2$	✓	✗	Γ_Z
$\mathcal{U}_5$	✓	✗	R_τ
$\mathcal{Q}_1$	✓	✗	R_τ
$\mathcal{Q}_5$	✓	✗	Γ_Z
$\mathcal{X}$	✓	✗	R_τ
$\mathcal{Y}_1$	✓	✗	Γ_Z
$\mathcal{Y}_5$	✓	✗	Γ_Z

$b \rightarrow s \ell \ell$ at FCC-ee

Bordone, Cornella, Davighi 2503.22635

- > Prospects from $B \rightarrow K^* \ell \ell$, $B_s \rightarrow \mu \mu$
- > Complementary to LHCb, CMS




An explicit (simplified) model

U_1 leptoquark + Z'

- > $SU(4)$ -unification inspired construction
- > Heavy $U_1 \sim (3, 1, 2/3)$ and Z' vectors with third-gen. coupling
- > $\mathcal{C}_{\ell q}^{(1)[3333]} = \mathcal{C}_{\ell q}^{(3)[3333]} = \frac{g_4^2 v^2}{8M_U^2}$
- > $\mathcal{C}_{\ell q}^{(1)[3333]} = -\frac{g_4^2 v^2}{32M_Z^2}$

[LA, Isidori, Pešut 2503.17019]




An explicit (simplified) model

U_1 leptoquark + Z'

- > $SU(4)$ -unification inspired construction
- > Heavy $U_1 \sim (3, 1, 2/3)$ and Z' vectors with third-gen. coupling
- > $\mathcal{C}_{\ell q}^{(1)[3333]} = \mathcal{C}_{\ell q}^{(3)[3333]} = \frac{g_4^2 v^2}{8M_U^2}$
- > $\mathcal{C}_{\ell q}^{(1)[3333]} = -\frac{g_4^2 v^2}{32M_Z^2}$

[LA, Isidori, Pešut 2503.17019]




> Central values and errors used in the FCC projections

Observable	SM	FCC projection
$B_c \rightarrow \tau \nu$	$(1.95 \pm 0.09) \times 10^{-2}$	$(2.09 \pm 0.03) \times 10^{-2}$
$B \rightarrow K \nu \bar{\nu}$	$(4.44 \pm 0.30) \times 10^{-6}$	$(5.64 \pm 0.17) \times 10^{-6}$
$B \rightarrow K^* \nu \bar{\nu}$	$(9.8 \pm 1.4) \times 10^{-6}$	$(12.4 \pm 0.4) \times 10^{-6}$
$B \rightarrow K \tau \tau$	$(1.64 \pm 0.06) \times 10^{-7}$	$(4.2 \pm 0.8) \times 10^{-6}$
$B_s \rightarrow \tau \tau$	$(7.45 \pm 0.26) \times 10^{-7}$	$(2.18 \pm 0.22) \times 10^{-5}$
$\Delta M_{B_s} / \Delta M_{B_s}^{\text{SM}}$	1.0	0.862 ± 0.015
$ g_\tau / g_\mu $	1.0	0.99926(7)
N_{eff}	3.0	2.9979(6)
A_τ	0.147	0.14668(21)
A_b	0.935	0.93502(22)

Vector-like fermions

- > $D \sim (3, 1, -1/3)$
- > Flavour: modified $Z\bar{b}_L s_L$ vertex
- > Sensitivity to ε_F (spurion)



- > $E \sim (1, 1, -1)$
- > Flavour: insensitive to M_E
- > EW: modified $Z\tau_L\tau_L$ vertex

