



Higgs boson pair production at LHC

Overview of the most recent results and HL
and FCC projections


Angela Taliencio

09/03/2026

Once upon a time



Once upon a time



Total luminosity collected is $\sim 474\text{ fb}^{-1}$ roughly a factor 100 more than what we had for the Higgs discovery!

What do we do with this enormous amount of data???

Once upon a time




Higgs properties

$$L_\phi = (D^\mu \phi)^\dagger (D_\mu \phi) - \sum_f g_f (\bar{\psi}_L \phi \psi_R + h.c.) - V(\phi)$$

Main Higgs properties:

- masses to vector bosons
 - from vev and the gauge couplings
- masses of the fermions
 - from Yukawa coupling



What is this plot telling us???


- SM is very solid...
- We are missing the **Higgs**

Introduction: HH production in the SM



Higgs SM potential

$$V(\phi^\dagger\phi) = \mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2$$

It is produced with



Destructive interference between the two diagrams



Very tiny cross section → very challenging analyses

$$\sigma_{SM}^{HH} = 31.05^{+5\%}_{-7\%} \text{ fb (scale } \oplus \text{ PDF } \oplus \alpha_S \oplus m_t)$$

Why is it that so important???

Introduction: HH production in the SM


In Standard Model destructive interference of triangle and box contributions \rightarrow tiny cross section \rightarrow Experimentally very challenging

$$V(\phi^\dagger\phi) = \mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2$$



The direct measure of λ is a strong test of the Electroweak Symmetry Breaking (EWSB) mechanism

Deep fundamental questions

- ❖ What is the order of the EW phase transition?
- ❖ What is the fate of the Universe? Is it stable?




Introduction: HH production in the SM





strong test of the Electroweak Symmetry Breaking (EWSB) mechanism

Deep fundamental questions

- ❖ What is the order of the EW phase transition?
- ❖ What is the fate of the Universe? Is it stable?



Introduction: HH production in the SM



- ❖ What is the order of the EW phase transition?
- ❖ What is the fate of the Universe? Is it stable?

Introduction: HH production in the SM


The direct measure of λ is a strong test of the SM prediction ($\lambda \sim 0.13$ from theoretical SM prediction)

$$V(\phi^\dagger\phi) = \mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2$$

- HH gluon-gluon fusion production (around 4000 events produced in the full RunII)

$$\sigma_{ggF}^{HH} = 31.05^{+2\%}_{-7\%} \text{ fb (scale } \oplus \text{ PDF } \oplus \alpha_S \oplus m_t)$$

*remark:
 κ_λ is a coupling modifier: $\kappa_\lambda = c_\lambda/c_{SM}$ Where c is the coupling




Gives a direct handle on the κ_λ parameter + κ_t dependence in the box diagram

Introduction: HH production in the SM


- HH gluon-gluon fusion production (around 4000 events produced in the full RunII)

$$\sigma_{ggF}^{HH} = 31.05^{+2\%}_{-7\%} \text{ fb (scale } \oplus \text{ PDF } \oplus \alpha_S \oplus m_t)$$

*remark:
 κ_X is a coupling modifier: $\kappa_X = c_X/c_{SM}$
 Where c is the coupling



Gives a direct handle on the κ_λ parameter + κ_t dependence in the box diagram




- Di Higgs kinematic strongly dependent on κ_λ
- Different m_{HH} ranges can be explored to be sensitive at different physics scenarios

Introduction: HH production in the SM

- HH VBF production (around 300 events produced in the full RunII)


$$\sigma_{VBF}^{HH} = 1.73 \pm 0.04 \text{ fb (scale } \oplus \text{ PDF } \oplus \alpha_S \oplus m_t)$$





VBF production mode is less dominant but gives a direct handle on the κ_V and κ_{2V} couplings

Introduction: HH production in the SM

HH gluon-gluon fusion production



HH VBF fusion production



- **Gluon fusion:** dominant production mode
- **Large destructive interference** \rightarrow tiny cross section

LHC Run2

CMS and ATLAS experiments

CMS DETECTOR

Total weight : 14,000 tonnes
 Overall diameter : 15.0 m
 Overall length : 28.7 m
 Magnetic field : 3.8 T

STEEL RETURN YOKE
 12,500 tonnes

SILICON TRACKERS
 Pixel ($100 \times 150 \mu\text{m}^2$) $\sim 1.9 \text{ m}^2 \sim 124\text{M}$ channels
 Microstrips ($80\text{--}180 \mu\text{m}$) $\sim 200 \text{ m}^2 \sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID
 Niobium titanium coil carrying $\sim 18,000 \text{ A}$



MUON CHAMBERS
 Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
 Endcaps: 540 Cathode Strip, 576 Resistive Plate Chambers

PRESHOWER
 Silicon strips $\sim 16 \text{ m}^2 \sim 137,000$ channels

FORWARD CALORIMETER
 Steel + Quartz fibres $\sim 2,000$ Channels



CRYSTAL ELECTROMAGNETIC CALORIMETER (ECAL)
 $\sim 76,000$ scintillating PbWO_4 crystals

HADRON CALORIMETER (HCAL)
 Brass + Plastic scintillator $\sim 7,000$ channels



HH searches in CMS

$H \rightarrow XX$





$bb\gamma\gamma$, $bbTT$, $4b$ most promising channels

HH production cross-section

Limit on the HH cross-section relative to its SM prediction (signal strength μ) is shown in those plots:

- 'Golden channels' are $4b$, $b\bar{b}\gamma\gamma$, $b\bar{b}\tau\tau$
- ATLAS combination leads to $2.9 \times SM$ (exp $2.4 \times SM$), CMS combination leads to $3.5 \times SM$ (exp $2.5 \times SM$)




- Also limits on the VBF production mode available, both experimet are sensitive to $\sim 90 \times SM$

CMS + ATLAS combination

Limit on the HH cross-section relative to its SM prediction (signal strength μ) is shown in those plots:

- 'Golden channels' are $4b$, $bby\gamma$, $bb\tau\tau$
- ATLAS combination leads to $2.9 \times \text{SM}$ (exp $2.4 \times \text{SM}$), CMS combination leads to $3.5 \times \text{SM}$ (exp $2.5 \times \text{SM}$)




Observed significance = 1.1
Expected significance = 1.3
for the SM HH signal

And this is 'only' with $\sim 138 \text{ fb}^{-1}$
at the end of the run we
expect at least 500 fb^{-1} per
experiment...

CMS + ATLAS combination


Both collaborations provided results for k_λ , k_τ and k_{2V} relative results

- k_λ scan dominated by $b\bar{b}\gamma\gamma$ and $b\bar{b}\tau\tau$
- k_{2V} scan dominated by $4b$ boosted signature




[CERN-EP-2026-011](#)

Constrain on the k_λ [-1.4, 6.6] obs - CMS
 Constrain on the k_λ [-1.3, 7.2] obs - ATLAS




Constrain on the k_{2V} [0.66, 1.4] obs - CMS
 Constrain on the k_{2V} [0.57, 1.5] obs - ATLAS
 $k_{2V} = 0$ excluded at $>6\sigma$ from both collaborations

bbyy in a nutshell



Tiny branching ratio $\sim 0.26\%$, still a golden channel

Easy to trigger, good mass resolution ($\sim 1-2\%$), high S/B






Triangle diagram dominates at low M_{HH}

\Rightarrow Excellent sensitivity to variations in κ_λ

bbyγ in a nutshell

Backgrounds:

- Non-resonant QCD: $\gamma\gamma$ +jets and γ +jets
- Single Higgs production



Mass resolution is extremely good, there are mainly two approaches to this analysis:


- Use the m_{jj} informations as input in a DNN and fit the $m_{\gamma\gamma}$ only (1D approach – ATLAS)
- DNN agnostic from m_{jj} , fit simultaneously $m_{\gamma\gamma}$, m_{jj} (2D approach – CMS)

bbyy analysis strategy: CMS

Two complementary approaches

- 1D approach:
 - multivariate BDT in which the m_{jj} information is explored at the ML level
 - Fit of $m_{\Upsilon\Upsilon}$
- 2D approach:
 - multivariate DNN, agnostic from m_{jj}
 - Fit simultaneously ($m_{\Upsilon\Upsilon}$ and m_{jj})

- **Boosted category** that complements both fitting approaches




Final results

bbyy analysis strategy: CMS

Two complementary approaches

- 1D approach:
 - multivariate BDT in which the m_{jj} information is explored at the ML level
 - Fit of $m_{\gamma\gamma}$
- 2D approach:
 - multivariate DNN, agnostic from m_{jj}
 - Fit simultaneously ($m_{\gamma\gamma}$ and m_{jj})
- **Boosted category** that complements both fitting approaches



Custom mbb regression implied in this channel → improvement of ~10% in the resolution

bbyy analysis strategy: classifiers

Single DNN model with 4 output classes:

- Nonresonant bkg (Data driven QCD + γ +jets, $\gamma\gamma$ +tt/jets)
- ttH
- Other single H bkg (ggH, bbH, VBFH, VH)
- Single: SM ggHH

Preprocessing:

- $m_{\gamma\gamma}/m_{jj}$ - decorrelation to allow for a 2D fit.

Architecture:

- 5 hidden layers, 1024 neurons each, ELU activation


bbyy analysis strategy: categorization

Signal Region optimization

- Maximise the asymptotic significance (Z) computed purely by MC samples + Data driven estimate

$$Z = \sqrt{2 \left[(s + b) \ln \left(1 + \frac{s}{b} \right) - s \right]}$$

- **Sequentially optimising** the SR boundaries, by choosing lower-cuts (upper-cuts) for the ggF HH score (to reject other processes)
 - Progressively add more SRs until the summed-in-quadrature significance plateaus



➔ Cumulative significance plateaus after 3 SRs



bbyy final fits

For the **resolved** topology we perform:

- Fit in the $m_{\gamma\gamma}$ leg:
 - **Non-resonant background:** family of smoothly falling functions + envelope method
 - **Resonant bkg + signal:** fitted with sum of gaussians
- Fit in the m_{bb} leg:
 - **Non-resonant background:** family of smoothly falling functions + envelope method
 - **ttH + VH + signal:** fitted with sum of gaussians
 - **VBFH + singleH:** fitted with an exponential



bby postfits

For the **resolved** topology we perform:




bbyy analysis strategy: boosted category

- The boosted Higgs tagging can significantly enhance significance, and it represents a **novelty** wrt previous CMS/ATLAS HH→bbyy results
- We investigated the potential sensitivity gains from having a separate boosted Higgs category
- “boosted aware” categorization procedure
 - Boosted category prioritized over resolved → (merged jet reconstruction of takes precedence over resolved reconstruction if an event can be reconstructed in both ways)



bbyy analysis strategy: results UL





- Sensitivity of the 3 signal region category is shown together with their combination
- Upper limit of **11 (7.3) x SM**

*bbgg Run2 (138 fb⁻¹) result 7.7 (5.2) obs (exp) xSM with >2x lumi, very promising results

*Largest source of systematic uncertainty is the HH theory uncertainty

bby γ analysis strategy: results κ_λ scan



[CMS-PAS-HIG-25-007](#)

Allowed κ_λ ranges

- Observed $\kappa_\lambda \in [-5.2, 12.0]$
- Expected $\kappa_\lambda \in [-3.6, 10.5]$

bbyy analysis strategy: summary

- With this early run3 results we achieved 7x SM with just half of the dataset implied in run2 → very promising result!!!

What's new????



Two complementary approaches explored at the same time


- 1D approach: multivariate BDT in which the m_{jj} information is explored at the ML level → Fit of $m_{\gamma\gamma}$
- 2D approach: multivariate DNN, agnostic from m_{jj} → Fit simultaneously ($m_{\gamma\gamma}$ and m_{jj})
- Both of them are complemented with a boosted category

Stay tuned!!! Many more to come!!!

bby analysis strategy: ATLAS

New kinematic likelihood fit procedure introduced to improve mbb resolution.

- Jet momentum varied within experimental resolution
- Exploits momentum balance against the well-reconstructed photons
- First iteration for unbiased estimate of mbb
- Second iteration with constrained mbb to recalculate other variables





46% improvement on mHH resolution

5% improvement on expected significance

bbyγ analysis strategy: ATLAS



- Separate BDTs trained in low and high mass regions
 - Input variables : kinematics of photons, b-jets, bb-system, missing ET, HT, 'single-topness'...
- Fit $m_{\gamma\gamma}$ distribution in 7 signal regions based on BDT score.



bbyy analysis strategy: CMS

- Observed signal strength: $\mu_{HH} = 0.9 +1.4/-1.1$
- Expected upper limit on μ_{HH} (assuming $\mu_{HH} = 0$): $2.6 \times SM$

CERN-EP-2025-140




Observed significance of SM HH: 0.8σ

Expected significance: 1.0σ

bbbb analysis in a nutshell


- Despite having the largest branching ratio, this channel is challenged by overwhelming QCD multijet backgrounds
- The QCD background modeled from simulation is limited and lacks higher-order corrections
- A data-driven background estimation becomes crucial for an accurate measurement

Resolved topology



- Separable H decay products (b jets)
- Reconstructed the Higgs candidates from identify 4 b-tag jets

Boosted topology





- Both the Higgs candidates are boosted
- Higgs candidates are reconstructed from two jets


bbbb analysis: data-driven bkg estimation

A multi-dimensional background reweighting technique, based on a DNN, is used

- Trained on the $CR_{2b} \rightarrow CR_{4b}$
- Applied to $SR_{2b} \rightarrow SR_{4b}$




bbbb analysis: results



- Combining the resolved & merged 22-23 analyses, we obtain an observed (expected) upper limit of **4.4 (4.4) × SM**, including all uncertainties.
- With 62 fb⁻¹, HH → 4b achieves a comparable sensitivity with Run 2 (138 fb⁻¹), and better than the Run 2 searches in bbγγ and bbττ

HL LHC: introduction

- 3x increase of peak luminosity
 - Data equivalent to 3000 fb⁻¹ in 10 years of operation
 - Upgraded CMS detector to cope with higher pileup and radiation damage




~ 36.7M HH events at
3000 fb⁻¹ for
CMS+ATLAS

~ 367M singleH events at
3000 fb⁻¹ for
CMS+ATLAS

Lots of statistics →
precise measurements

HL LHC: introduction


- 3x increase of peak luminosity
 - Data equivalent to 3000 fb⁻¹ in 10 years of operation
 - Upgraded CMS detector to cope with higher pileup and radiation damage



- European Strategy for Particle Physics is the cornerstone for Europe's long-term decision making process
- Last update from 2020
- Priority on the successful completion of the High-Luminosity LHC over the coming decade
- Higgs factory as the highest priority to follow the LHC

2025 ESU update - updated HL-LHC projections


- We want to emphasize the unique opportunity that HL will give:
 - more lumi + new analysis technique explored in the past 5 years unique update the expected precision on the Higgs couplings



2025 ESU update - updated HL-LHC projections

- We want to emphasize the unique opportunity that HL will give:
 - update the expected precision on the Higgs couplings
 - establishing the shape of the electroweak vacuum
 - stability of the universe

First attempt to give a proper answer to those questions



HH results - Significance

- 3000 fb⁻¹ per experiment, run3 object improvement included

| Channel | HH Significance ATLAS | HH Significance CMS |
|--------------------|-----------------------|---------------------|
| bb $\tau\tau$ | 3.8 | 2.7 |
| bb $\gamma\gamma$ | 2.6 | 2.6 |
| 4b resolved | 1.0 | 1.3 |
| 4b boosted | - | 2.2 |
| Multilepton | 1.0 | - |
| bb $\ell\ell$ | 0.5 | - |
| Combination | 4.5 | 4.5 |
| ATLAS + CMS | 7.60 | |

- Evidence with single experiment
- **Combined evidence >7 σ**

[ATL-PHYS-PUB-2025-018 CMS-HIG-25-002](#)

HH results - κ_λ precision


- 3000 fb⁻¹ per experiment, run3 object improvement included

| Channel | κ_3 precision 68% CL | κ_3 precision 68% CL |
|--------------------|-----------------------------|-----------------------------|
| bb $\tau\tau$ | [0.5, 1.6] | [0.3, 2.0] |
| bb $\gamma\gamma$ | [0.5, 1.7] | [0.4, 1.9] |
| 4b resolved | [-0.5, 6.1] | [-0.3, 7.2] |
| 4b boosted | - | [-0.4, 8.2] |
| Multilepton | [-0.1, 4.7] | - |
| bb $\ell\ell$ | [-2.1, 9.1] | - |
| Combination | [0.6, 1.4] | [0.6, 1.5] |
| ATLAS + CMS | -26/+29 | |

- ~26% precision for $\kappa_\lambda=1$


[ATL-PHYS-PUB-2025-018 CMS-HIG-25-002](#)

HH results - κ_λ precision



[ATL-PHYS-PUB-2025-018 CMS-HIG-25-002](#)

- κ_1 scan assuming different values of κ_1
 - maximum interference between box and triangle diagram at 2.45 \rightarrow largest precision




Shape of the electroweak vacuum

- Measuring the Higgs self-coupling alone is insufficient to fully determine or constrain the shape of the EWSB potential
- To provide model-independent conclusions on fundamental questions, such as whether the EWSB transition is first-order (a key element for electroweak baryogenesis) **an assumption about the form of the potential is generally required:**
 - dim-6 EFT
 - dim-8 EFT
 - logarithmic
 - exponential

Shape of the electroweak vacuum


- The light blue delimits the area where a first order phase transition would be possible in a dimension 6 EFT.
 - 2 σ full exclusion is reached at 3 ab⁻¹ for the EFT dim 6 FOPT inducing potentials



[ATL-PHYS-PUB-2025-018 CMS-HIG-25-002](#)

What's next????

FCC-project




FCC-hh

pp collider

FCC-hh has three baselines:

- 84 TeV (14T Magnet)
- 100 TeV (17T Magnet)
- 120 TeV (20T Magnet)



| Project | IP | Higher Energy |
|---------|----|---|
| FCC-hh | 2 | 84.6 TeV: 30 ab^{-1} 100 TeV: 30 ab^{-1} 120 TeV: 12 ab^{-1} |

Re-optimized $\bar{b}b\gamma\gamma$ analysis strategy

$$m_x = m_{\bar{b}b\gamma\gamma} - m_{\bar{b}b} - m_{\gamma\gamma} + 250\text{GeV}$$

Pre-selected events




$m_x < 350$
GeV

$m_x > 350$
GeV





DNNs

categorization to
suppress the bkg



- Shape depends on κ_λ
- Region $m_x < 350 \text{ GeV}$ has low S/B for $\kappa_\lambda=1$ (SM), but contributions from $\kappa_\lambda \neq 1$ (BSM) signals

Re-optimized $\bar{b}b\gamma\gamma$ analysis strategy



Re-optimized $\bar{b}b\gamma\gamma$ analysis strategy



Pre-selected events



DNNs categorization to suppress the bkg





Categorize based on $m_{\bar{b}b}$





Fit $m_{\gamma\gamma}$

FCC-hh new results

- Precision on k_l at 84 TeV (st only and syst scenarios) on the left
- Precision on k_l as a function of the the m_{jj} resolution on the right




Conclusions



- ATLAS and CMS program on diHiggs is extremely exciting and is performing behind expectations
 - Looking forward for the run3 combination!
- Many more to come!!**

Backup

CMS reconstruction



EFT Frameworks used in HH

- The **kappa-framework**: Modifies the couplings predicted by the SM

$$\kappa_\lambda = \frac{\lambda}{\lambda_{SM}}$$



- Higgs EFT (HEFT)**: **Natural extension** of the **kappa-framework** with additional BSM couplings
- Standard Model EFT (SMEFT)**: Similar to HEFT but the **Higgs is a SM doublet** adding correlations between H and HH

| | ATLAS | | CMS | |
|-----------------|-------|-----|-----|-----|
| HH production | ggF | VBF | ggF | VBF |
| kappa-framework | ✓ | ✓ | ✓ | ✓ |
| HEFT | ✓ | | ✓ | |
| SMEFT | ✓ | ✗ | ✗ | ✗ |



HEFT results

Both collaborations provided results in the HEFT framework:

- **CMS new results:** c_2 scan dominated by the $b\bar{b}\tau^+\tau^-$ channel
- **ATLAS 2D scan:** c_2 VS C_{gghh}




[Phys. Rev. Lett. 133 \(2024\) 101801](#) C_{gghh}




Higgs self-coupling and quartic HHVV coupling

Both collaborations provided results for k_λ , k_t and k_{2V} relative results

- k_λ scan dominated by $b\bar{b}\gamma\gamma$ and $b\bar{b}\tau\tau$
- k_{2V} scan dominated by $4b$ boosted signature



Constrain on the k_λ [-1.4, 6.4] obs - CMS
 Constrain on the k_λ [-1.2, 7.2] obs - ATLAS




Constrain on the k_{2V} [0.6, 1.4] obs - CMS
 Constrain on the k_{2V} [0.6, 1.5] obs - ATLAS
 $k_{2V} = 0$ excluded at $>6\sigma$ from both collaborations

bbyy analysis strategy: classifiers

- Major backgrounds categorized into three different classes based on their similarity to signal kinematics:

- Non-resonant QCD, $\gamma(\gamma)$ +jets, ggH , VBF H
- ttH & bbH - has true b-jets
- $V(bb)H$ - has m_{bb} resonance



- Single DNN model with 4 output classes:
 - Nonresonant bkg (Data driven QCD + γ +jets, $\gamma\gamma$ + tt /jets)
 - ttH
 - Other single H bkg (ggH , bbH , VBFH, VH)
 - Single: SM $ggHH$
- Preprocessing:
 - $m_{\gamma\gamma}/m_{jj}$ - decorrelation to allow for a 2D fit.
- Architecture:
 - 5 hidden layers, 1024 neurons each, ELU activation

bbyy analysis strategy: categorization

- The Multiclass BDT provides 4 outputs: $P_{\text{signal}}, P_{\text{t\bar{t}H}}, P_{\text{VH}}, P_{\text{QCD}}$
- We define two discriminators:

$$D_{\text{t\bar{t}H}} = \frac{P_{\text{HH}}}{P_{\text{HH}} + P_{\text{t\bar{t}H}}} \quad D_{\text{QCD}} = \frac{P_{\text{HH}}}{P_{\text{HH}} + P_{\text{QCD}} + P_{\text{VH}}}$$


- Optimize cuts on D_{QCD} and $D_{\text{t\bar{t}H}}$ to maximize S/B
 - Start with $D_{\text{t\bar{t}H}}$ cut to achieve 85% t\bar{t}H rejection.
 - Scan D_{QCD} and find max of S/B
- 3 SR category are selected
- Fit is performed on the m_{VY} leg

Signal Region optimization



- Maximise the asymptotic significance (Z) computed purely by MC samples + Data driven estimate

$$Z = \sqrt{2 \left[(s + b) \ln \left(1 + \frac{s}{b} \right) - s \right]}$$

- **Sequentially optimising the SR** boundaries, by choosing lower-cuts (upper-cuts) for the ggF HH score (to reject other processes)
 - Progressively add more SRs until the summed-in-quadrature significance plateaus




bbyy analysis strategy: results UL



*Largest source of systematic uncertainty is the HH theory uncertainty

*bbgg Run2 (138 fb⁻¹)
result 7.7 (5.2) obs
(exp) xSM

bby γ analysis strategy: results κ_λ scan



Allowed κ_λ ranges

- Observed $\kappa_\lambda \in [-3.9, 10.4]$
- Expected $\kappa_\lambda \in [-4.5, 11.1]$

HH projections: k_λ precision and significance

CMS and **ATLAS** provided projections for HL:

- evidence of HH production at the end of HL for both experiments

