



Overview of 21 cm Experiments




Peter Timbie
Department of Physics
UW-Madison

8-10 Sept 2015

2015 Tianlai Colla
21 cm Intensity M



21 cm cosmology timeline







Pritchard & Loeb Rep. Prog. Phys. 75 (2012)


21 cm Cosmology Goals

- **Observe global 21-cm spectrum**
EDGES, SCI-HI, DARE
- **Observe the Cosmic Dark Ages ($z > 10$)**
LWA
- **Observe Epoch of Reionization ($z \sim 10$)**
CMA21, PAPER, MWA, LOFAR
- **Observe Large-Scale Structure at ‘Late-times’ ($z < 3$)**
dark energy
non-Gaussianity (probe of inflation)
neutrino mass, etc.
BINGO, FAST, **GBT, Parkes, CHIME, HIRAX, PAON, TIANLAI, SKA**
- **Monitor time-dependent sky**
Pulsars, FRB's

BAO: Baryon Acoustic Oscillations in pre-recombination era → CMB acoustic peaks & a standard ruler in the galaxy density field




HI and Optical Surveys Detect Similar Structure to $z \sim 0.07$



Haynes et al. (2011)




21 cm surveys can explore huge comoving volumes



Galactic Foregrounds are $\sim 10^4$ X Brighter than HI at $z \sim 1$!

de Oliveira-Costa 'GSM' Model (2008)



First Steps with Intensity Mapping at GBT




- clean beam, unblocked aperture (100 m)
- quiet cooled receiver ($T_{\text{sys}} \sim 25$ K)
- for 700–900 MHz band ($z \sim 0.59 - 1.0$)
- angular resolution good for intensity mapping
 $\sim 0.25^\circ \rightarrow 10 h^{-1} \text{ Mpc}$ at $z \sim 0.8$
- available for ~ 100 's of hours
- ASIAA, CMU, CITA (+ NAOC, UW-Madison)

Tests of:


- foreground removal
- RFI removal
- HI intensity mapping
 - cross-correlation with optical surveys
 - autocorrelation
- HI abundance (Ω_{HI})

WiggleZ Dark Energy Survey: correlate with GBT survey


Drinkwater et al. MNRAS (2010)



- large-scale spectroscopic survey
- emission-line galaxies
- selected from UV and optical imaging
- ~ 350 galaxies deg^{-2}
- 1000 deg^2
- Anglo-Australian telescope



RFI Jungle



Mon Jun 24 13:24:58 2013


Removal techniques

- High resolution spectra (200 MHz/ 4096 ~ 50 kHz bins)
- Variance
- Linear polarization

E. Switzer



GBT map before foreground removal WiggleZ field

Masui et al. ApJ 763:L20 (2013)



Foreground removal

- *Instrumental effects* (polarization leakage from $P \rightarrow I$) and Faraday rotation produce structure in spectra of bright foregrounds.
- In each angular pixel, use **singular value decomposition (SVD)** to find largest modes in frequency (redshift) direction.
- Simulate/ correct for signal subtracted by this method.
- In *cross-correlation* with galaxy survey, unremoved foregrounds increase noise, but don't bias the power spectrum. Optimum number of cut modes ~ 15 .
- In *auto-correlation*, unremoved foregrounds bias the power spectrum.




Switzer et al. 1504.07527v1

Cleaned 21-cm GBT map

Masui et al. ApJ 763:L20 (2013)


GBT 15hr field, cleaned, beam convolved (800.4 MHz, $z = 0.775$)



$$\frac{T_{\text{Sys}}}{\sqrt{\Delta\nu\tau}} = \frac{26}{\sqrt{0.78 \text{ MHz} \times 2200 \text{ s}}} = 0.63 \text{ mK/beam}$$


HI – WiggleZ cross-power spectrum

Masui et al. ApJ 763:L20 (2013)





Fast Radio Burst Discovered!

Pulse arrival time vs frequency for FRB 110523




Masui et al. 2015

GBT 800 MHz 7-element array




Each feed antenna has an independent cryostat & cooler

ASIAA, NRAO, UW-Madison



Parkes 21 cm Survey



- ASIAA, CITA, CMU, NAOC, UW-Madison
- 0.25° beam $\sim 1.5 h^{-1}$ Mpc
- cross-correlate with 2dF galaxy redshift survey
 - 220,000 galaxies, $z \sim 0.11$
- 300 hrs being analyzed now
 - $1500 \text{ deg}^2 \sim$ most of 2dF volume
 - $10^7 h^{-3} \text{ Mpc}^3$
- parallel mapmaker for multibeam receiver





Selected HI Large-Scale Structure Instruments

		T_{inst}	$N_d \times N_b$	D_{dish} (m)	z_{min}	z_{max}	S (deg ²)
single dish	GBT	29	1x1(7)	100	0.54	1.09	100
	Parkes	23	1x13	64	0.00	0.23	5,000
	<i>BINGO</i>	50	1x50	25	0.13	0.48	5,000
	<i>FAST</i>	20	1x20	500	0.42	2.55	2,000
interferometer	CHIME 4, 100 m cyl.	50	1280x1	20	0.77	2.55	25,000
	Tianlai 8, 120 m cyl.	50	2048x1	15	0.00	2.55	25,000
	HIRAX	50	1024x1	5	0.77	2.55	15,000
interf. + autocor.	<i>SKA1-MID</i>	20	190x1	15	0.00	3.06	25,000
	<i>SKA1-SUR</i>	30	96x36	15	0.23	1.38	25,000

from Bull *et al.* ApJ 803 (2015)

BAO forecasts




$$\frac{p}{\rho} = w(a) \sim w_0 + \frac{z}{1+z} w_a$$


Bull et al. (2015)

BINGO

BAO from Integrated Neutral Gas Observations




- University of Manchester
- Battye et al. MNRAS 434 (2013)
- single-dish using natural landscape
- drift scans
- map-making on large scales uses reference beams to view NCP or SCP



Bull et al. 2015

PAON-4

Paraboles A l'Observatory de Nancay



Université Paris-Sud
Observatoire de Paris
Laboratoire Lagrange
Centre d'Etudes de Saclay

- 5-m diameter dishes in transit array
- FPGA – based BAORadio backend
- tests of 21-cm intensity mapping




1st light ceremony April '15

HIRAX

Hydrogen Intensity and Real-time Analysis eXperiment



UKZN, DUT South Africa
SKA site Karoo desert
1024, 6 meter dishes (1st from Beifang)
close-packed regular array
observations start 2018
CHIME correlator
other science:
FRB, pulsars, HI absorbers



CHIME

(Canadian Hydrogen Intensity Mapping Experiment)

Prototype

2, 8m dishes
4 channels


Pathfinder

2, 20mx40m cylinders
256 channels, 400 MHz

Full CHIME - funded


4, 20mx100m cylinders
1280 channels, 400 MHz
operating in 2016

UBC
DRAO
McGill
NRC-CNRC
Toronto




CHIME calibration

Beam:
pulsar 'holography':
correlate cylinder feeds
with 26 m dish



Gain:
broadband noise
injection



Laura Newburgh 1406.2267

Summary: big steps soon for HI ‘tomography’

In the next 12 months Tianlai and other 21-cm intensity-mapping experiments will have interferometer sky data:

- first light for Tianlai
- tests of correlators
- calibration
- maps
- foreground removal from large arrays

