

# Algal bioconvection in confined dispersed media

*lundi 17 juin 2024 15:40 (20 minutes)*

Active particles are inherently out-of-equilibrium systems, able to uptake energy from their environment and convert it to motion.

For example, *Chlamydomonas Reinhardtii* (CR) is a micro-swimmer whose orientation can be dictated by a light gradient in its environment (phototaxis). It is known that a collective motion triggered due to the phototaxis of a population of CR generates a nonlinear phenomenon: bioconvective structures that affect the fluid medium [1].

The purpose of this experimental study is to control the motion of algae swarms and resulted bioconvective vortices in order to achieve a guided transportation of microscopic objects submerged in the algal suspension. High concentrations of algae and microparticles were confined in a small square Hele-Shaw cell surrounded by a series of LED, allowing us to apply different well-controlled gradients of light stimulus in a quasi-2D horizontal domain.

It was shown that the microparticles can be transported to a target zone by controlling the displacement of algal bioconvective structures.

[1] J. Dervaux, M. Capellazzi Resta, and P. Brunet, *Nature Phys* 13, 306–312 (2017).

**Auteur principal:** NICOLAZO-CRACH, Victoria (Laboratoire FAST)

**Co-auteurs:** M. LAROUSSI, Taha (LadHyx); M. BOUVARD, Julien (LadHyx); Dr AMSELEM, Gabriel (LadHyx); JARRAHI, Mojtaba (université Paris-Sud)

**Orateur:** NICOLAZO-CRACH, Victoria (Laboratoire FAST)

**Classification de Session:** Présentations