

Higgs interference effects at NLO


Raoul Röntsch

KARLSRUHE INSTITUTE OF TECHNOLOGY

Higgs Hunting 2016

Narrow width approximation

NWA: Factorize production and decay



Accuracy : $\Gamma_H/m_H \sim 10^{-5}$



BUT: 10% of events in $H \rightarrow VV$ above $2m_V$ threshold

(Kauer, Passarino '12)

➡ NWA **not sufficient** to describe behavior of $H \rightarrow VV$

Beyond the NWA

Consistent treatment requires both **signal** and **background** amplitudes




“signal” A_H

“background” A_b

$$|A_{ZZ}|^2 = |A_H|^2 + |A_b|^2 + 2\text{Re}[A_H A_b^*]$$

$$\rightarrow \sigma_{\text{full}} = \sigma_{\text{sigl}} + \sigma_{\text{bkgd}} + \sigma_{\text{intf}}$$

Interference effect and line shape




Interference is **strong** and **destructive**, especially at high invariant mass

Campbell, Ellis, Williams '13

Understanding high energy behavior

Cut open top loop – have $t\bar{t} \rightarrow VV$




J. Campbell

- **Signal** and **background** amplitudes-squared grow like E^2
- **Interference** grows like $-E^2 \rightarrow$ cancels E^2 terms of signal and background
- **Higgs unitarizing massive scattering amplitude** – connected to its role in EWSB and mass generation

Using off-shell Higgs

Off-shell Higgs events provide **a new tool** for studying Higgs physics

- High mass behavior probes **unitarization properties** of Higgs
 - **As important** as measuring Higgs couplings for understanding EWSB
 - Cross sections small \rightarrow HL-LHC
- Higgs couplings **more sensitive to NP** at high energies



Campbell, Ellis, Williams '13

Using off-shell Higgs: indirect width constraints

- Observation of Caola & Melnikov:

$$\sigma_{\text{on}} \propto g_i^2 g_f^2 / \Gamma_H \quad \sigma_{\text{off}} \propto g_i^2 g_f^2$$

$$\Rightarrow \Gamma_H \propto \frac{\sigma_{\text{off}}}{\sigma_{\text{on}}} \quad \longrightarrow \quad \text{indirect constraint on width}$$

- CMS: $\Gamma_H < 13 \text{ MeV}$ ATLAS: $\Gamma_H < 23 \text{ MeV}$
- Direct constraints $\sim 1 \text{ GeV}$
- Compare with SM value: $\Gamma_H \approx 4 \text{ MeV}$

Indirect Higgs width constraint: *caveat emptor*

Indirect constraints not model-independent:

- **Assume same couplings on- & off-shell**
- Can construct models with $\Gamma_H > \Gamma_H^{\text{SM}}$ but no sensitivity from off-shell measurements (Englert, Spannowksy '14)
- Possible option:
 - Introduce energy-dependent couplings in κ -framework / EFT (Englert, Soreq, Spannowsky '14)
 - Constrain couplings and width simultaneously
 - Highly non-trivial dependence of **signal**, **background** and **interference** on these couplings! (see e.g. Azatov, Grojean, Paul, Salvioni '16)

Impact of higher order corrections


- Signal (incl. top mass effects) known at **NLO**
 - **NLO corrections large**, k-factor ~ 1.7
- Experimental analyses use background and interference at **LO only**
 - NLO corrections approximated by corrections from signal
→ **adds uncertainties to analysis**
- Recent work has extended background and interference to **NLO**

Campbell, Czakon, Ellis, Kirchner, hep-ph/1605.01380


Caola, Dowling, Melnikov, R.R., Tancredi, hep-ph.1605.04610

Higgs Interference Effects at NLO

Known




Known




Massless: known

Massive: extremely difficult / impossible




Massless: easy


Massive: moderate



Known




Known



Spira, Djouadi, Graudenz, Zerwas '95; Harlander, Kant '05; Aglietti, Bonciani, Degrassi, Vicini '07;
 Ellis, Hinchliffe, Soldate, v.d. Bij '88;
 Caola et al '15, v. Manteuffel, Tancredi '15
 Hagiwara, Kuruma, Yamada '91; Campbell, Ellis, Zanderighi '07;
 v.d. Bij, Glover '89;

$gg \rightarrow (H) \rightarrow ZZ$: Top Mass Expansion

Expand in s/m_t^2




- Keep terms to $(s/m_t^2)^4$
- Expect to be valid for partonic energies

$$s \lesssim 4m_t^2$$

Dowling, Melnikov '15

Validity of top mass expansion

Can check validity at LO:



→ Good approximation below $2m_t$ threshold

Restricted to

$$m_{4\ell} \leq 2m_t \quad p_{T,j} < 150 \text{ GeV}$$

- Cannot probe unitarization effects :(
- Large window $150 \text{ GeV} \lesssim m_{4\ell} \leq 2m_t$ where Higgs is off-shell and we can study **interference effects at NLO**

Parameters

- $gg \rightarrow ZZ \rightarrow e^+e^-\mu^+\mu^-$ at 13 TeV LHC
- Dynamical scale $\mu_F = \mu_R = \{m_{4\ell}/4, m_{4\ell}/2, m_{4\ell}\}$
- Minimal cuts:
 - $150 \text{ GeV} \leq m_{4\ell} \leq 340 \text{ GeV}$
 - $p_{T,j} < 150 \text{ GeV}$
 - $60 \text{ GeV} \leq m_{\ell\ell} \leq 120 \text{ GeV}$

$gg \rightarrow (H) \rightarrow ZZ$ Results: Cross Sections

$$\sigma_{\text{LO}}^{\text{signal}} = 0.043_{-0.009}^{+0.012} \text{ fb}, \quad \sigma_{\text{NLO}}^{\text{signal}} = 0.074_{-0.008}^{+0.008} \text{ fb}$$





$$\sigma_{\text{LO}}^{\text{bkgd}} = 2.90_{-0.58}^{+0.77} \text{ fb}, \quad \sigma_{\text{NLO}}^{\text{bkgd}} = 4.49_{-0.38}^{+0.34} \text{ fb}$$

$$\sigma_{\text{LO}}^{\text{intf}} = -0.154_{-0.04}^{+0.031} \text{ fb}, \quad \sigma_{\text{NLO}}^{\text{intf}} = -0.287_{-0.037}^{+0.031} \text{ fb}$$

$$\sigma_{\text{LO}}^{\text{full}} = 2.79_{-0.56}^{+0.74} \text{ fb}, \quad \sigma_{\text{NLO}}^{\text{full}} = 4.27_{-0.35}^{+0.32} \text{ fb},$$


- **Destructive interference** ~ **5%**
 - ~ 4 x larger than signal, order of magnitude smaller than background
 - Can use specialized cuts needed to enhance relative to signal and background
- Scale uncertainty: **20%-30% at LO**, **10% at NLO**
- $K_{\text{sigl}} = 1.72$ $K_{\text{bkgd}} = 1.55$ $K_{\text{intf}} = 1.65 \simeq \sqrt{K_{\text{sigl}} K_{\text{bkgd}}}$

$gg \rightarrow (H) \rightarrow ZZ$ Results: Mass distributions




- Differential k-factors **relatively flat**...
- Except for interference near $2m_Z$ threshold

$gg \rightarrow (H) \rightarrow ZZ$ Results: Differential k-factor



K_{intf}



- **Massless loop** dominates near $2m_Z$ threshold, **drives k-factor behavior**

Comparison with similar work


Campbell, Czakon, Ellis, Kirchner, hep-ph/1605.01380

- Only **interference** contribution considered
- On-shell Z bosons, so $m_{ZZ} > 2m_Z$
- Massive two-loop amplitudes computed in mass expansion to $(s/m_t^2)^6$
- Massive real emission amplitudes computed exactly – no need for jet cut
- **Results extended beyond $2m_t$ threshold using Padé approximations – look at high-mass tail**

Comparison with similar work

Campbell, Czakon, Ellis,
Kirchner, arXiv:1605.01380


Qualitatively similar
behavior of k-factors
near $2m_Z$ threshold



k-factor flat
in high-
energy tail

$gg \rightarrow (H) \rightarrow WW$

- Analogous to $gg \rightarrow (H) \rightarrow ZZ$
- Mass expansion more complicated since **top and bottom quarks mix** in loop
- \rightarrow neglect 3rd generation altogether
 - **Comparable** to massless contribution at low-intermediate $m_{T,WW}$
 - **Dominate** at high $m_{T,WW}$
- **Partial results only**







- $gg \rightarrow W^+W^- \rightarrow \nu_e e^+ \mu^- \bar{\nu}_\mu$
- No kinematic cuts imposed
- Scales as for ZZ

$gg \rightarrow (H) \rightarrow WW$ Results: Cross Sections

$$\begin{aligned} \sigma_{\text{LO}}^{\text{signal}} &= 48.3^{+10.4}_{-8.4} \text{ fb}, & \sigma_{\text{NLO}}^{\text{signal}} &= 81.0^{+10.5}_{-8.2} \text{ fb} \\ \sigma_{\text{LO}}^{\text{bkgd}} &= 49.0^{+12.8}_{-9.7} \text{ fb}, & \sigma_{\text{NLO}}^{\text{bkgd}} &= 74.7^{+5.5}_{-6.2} \text{ fb} \\ \sigma_{\text{LO}}^{\text{intf}} &= -2.24^{+0.44}_{-0.59} \text{ fb}, & \sigma_{\text{NLO}}^{\text{intf}} &= -4.15^{+0.47}_{-0.54} \text{ fb} \\ \sigma_{\text{LO}}^{\text{full}} &= 95.0^{+22.6}_{-17.6} \text{ fb}, & \sigma_{\text{NLO}}^{\text{full}} &= 151.6^{+15.4}_{-13.9} \text{ fb}. \end{aligned}$$


- **Destructive interference** ~ 2%
 - Higgs peak present → interference smaller than signal and background
- Scale uncertainty reduced by factor ~ 2
- $K_{\text{sigl}} = 1.68$ $K_{\text{bkgd}} = 1.53$ $K_{\text{intf}} = 1.85$
 - slightly above geometric mean

$gg \rightarrow (H) \rightarrow WW$ Results: Mass distributions



- Differential k-factors **relatively flat...**
- ... except for interference near $2m_W$ threshold – as in ZZ case


$gg \rightarrow (H) \rightarrow WW$ Results: Estimating effect of 3rd generation



- As in ZZ case, enhancement from **massless loops**
- **3rd generation loops** give relatively flat differential k-factor

→ estimate by using LO results scaled by approximate k-factor

$$\sqrt{K_{\text{sigl}} K_{\text{bkgd}}}$$



Conclusions

- Higgs off-shell behavior provides a rich environment to study Higgs physics:
 - Probe **unitarizing behavior** of the Higgs
 - **Indirect constraints** on Higgs width
 - **Test couplings** at high energies
- NLO corrections to **interference** in $gg \rightarrow ZZ$ and $gg \rightarrow WW$, are now known, at least below $2m_t$ threshold
- Difficulty of computing two-loop massive corrections
 - **top mass expansion** for ZZ
 - **neglect 3rd generation** for WW
- ZZ in window $150 \text{ GeV} \leq m_{4\ell} \leq 340 \text{ GeV}$
 - Moderate k-factors ~ 1.6 - 1.7
 - $K_{\text{intf}} \simeq \sqrt{K_{\text{sigl}} K_{\text{bkgd}}}$ **except near $2m_z$ threshold** – driven by massless amplitudes
- WW :
 - Interference k-factor slightly larger than signal and background k-factors
 - Effect of 3rd generation at NLO approximated assuming uniform contribution to k-factor

THANK YOU!