

Séminaires doctorants 2ème année

May 10, 2021/Orsay

Two-pion exchange contributions to NN interaction in covariant baryon chiral perturbation theory

Yang Xiao (肖杨)


Collaborators:

U. van Kolck, Li-Sheng Geng,

Chun-Xuan Wang & Jun-Xu Lu.




Nuclear forces (NF) – Basic input in nuclear physics



Nuclear force from QCD



➤ Color confinement

Quark d.o.f.'s instead of hadrons



quarks

hadrons



➤ Asymptotic freedom

Coupling constants $\alpha_s \geq 1$ (low energy)

● Nonperturbative - unsolvable



- Lattice QCD
- **Effective Field Theory (we prefer)**
- Phenomenological models

Why Chiral Effective Field Theory

□ Chiral Effective field theory


- Low energy QCD
 - $L_{QCD} \rightarrow L_{\chi EFT} \sim \sum_{\mathbf{v}} c_{\mathbf{v}} \times L_{\chi EFT}^{\mathbf{v}}$ ($c_{\mathbf{v}}$ low energy constants, \mathbf{v} chiral order)
- Degree of freedom
 - quarks \rightarrow hadrons
- Power counting scheme (expansion parameter)
 - $Q \sim M_{lo}/M_{hi}$ ($M_{hi} \sim \Lambda_{\chi} \sim 1\text{GeV}$, $M_{lo} \sim p, m_{\pi}$)



□ Main advantages of Chiral Nuclear force

- Self-consistently include many-body forces
$$V = V_{2N} + V_{3N} + V_{4N} + \dots$$
- Systematically improve NF order by order
$$V_{iN} = V_{iN}^{LO} + V_{iN}^{NLO} + V_{iN}^{NNLO} + \dots$$
- Systematically estimate theoretical uncertainties

$$|V_{iN}^{LO}| > |V_{iN}^{NLO}| > |V_{iN}^{NNLO}| > \dots$$



Chiral NN potential is of high precision

Phenomenological forces

NR Chiral nuclear force

	Reid93	CD-Bonn	LO	NLO	N2LO	N3LO	N4LO
No. of para.	50	38	2	9	9	24	24
χ^2 /datum	1.03	1.02	94	36.7	5.28	1.27	1.10


D.Entem, et al., PRC96(2017)024004

Why covariant ?

- **Relativistic** approaches **successful**
 - **Atomic and molecular systems**, why gold is **yellow**
 - **Nuclear system**, spin-orbit splitting, pseudospin symmetry
 - **One-baryon sector**, magnetic moments, masses, sigma terms
- **A critical input** to nuclear reaction/structure calculations in **covariant framework**
 - **Dirac Brueckner Hartree-Fock**
 - **Covariant Chiral MF**
 -

LO covariant Chiral NN force

Xiu-Lei Ren, et.al. CPC42 (2018) 1, 014103









- **LO** relativistic \approx **NLO** non-relativistic in S & P wave !
- **Higher order** needs to be explored.
- **Our purpose**
 - ✓ Providing **key inputs** to high precision **covariant chiral nuclear force**

Higher order Feynman diagrams

2N Force


3N Force

4N Force



Key inputs

- Four nucleon (baryon) vertices
- Pion-nucleon vertices
- **Two-pion exchanges (TPE)**



Two-pion exchange up to $O(p^3)$


□ Covariant chiral Lagrangians:

$$\mathcal{L}_{\pi N}^{(1)} = \bar{N} \left(i\not{D} - m + \frac{g_A}{2} \not{\psi} \gamma_5 \right) N,$$


$$\mathcal{L}_{\pi N}^{(2)} = c_1 \langle \chi_+ \rangle \bar{N} N - \frac{c_2}{4m^2} \langle u^\mu u^\nu \rangle (\bar{N} D_\mu D_\nu N + h.c.) + \frac{c_3}{2} \langle u^2 \rangle \bar{N} N - \frac{c_4}{4} \bar{N} \gamma^\mu \gamma^\nu [u_\mu, u_\nu] N,$$

□ Feynman diagrams

$(Q/\Lambda)^2$



$(Q/\Lambda)^3$



Covariant chiral potentials

$$V_{NN}^{(2)} = \bar{u}_1 \bar{u}_2 \left\{ \begin{array}{c} \text{(a)} \\ \text{(b)} \\ \text{(c)} \\ \text{(d)} \\ \text{(e)} \end{array} \right\} u_1 u_2$$

$$V_{NN}^{(3)} = \bar{u}_1 \bar{u}_2 \left\{ \begin{array}{c} \text{(a)} \\ \text{(b)} \\ \text{(c)} \\ \text{(d)} \end{array} \right\} u_1 u_2$$

$$u(\mathbf{p}, s) = N \left(\frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{E + m_n} \right) \chi_s, \quad N = \sqrt{\frac{E + m_n}{m_n}}$$

T matrix & phase shifts

- On-shell T matrix: in first order perturbation theory

$$T_{NN} = V_{NN}$$

- T matrix to phase shifts


$$\delta_{LSJ} = -\frac{m_n^2 |\mathbf{p}|}{16\pi^2 E} \operatorname{Re} \langle LSJ | \mathcal{T}_{NN} | LSJ \rangle,$$

$$\epsilon_J = \frac{m_n^2 |\mathbf{p}|}{16\pi^2 E} \operatorname{Re} \langle J - 1, 1, J | \mathcal{T}_{NN} | J + 1, 1, J \rangle.$$



Perfect check of relativistic corrections (No free parameters)


Phase shifts-D wave



*Xiao et al.,
PRC 102, 054001 (2020)
Kaiser et al.
NPA 625, 758 (1997).*

- **Red bands:**
renormalization scale $0.5 \sim 1.5\text{GeV}$
- ✓ **Quantitatively better**
- ✓ ϵ_2 much improved
- **Contact terms** needed


Phase shifts-F wave



✓ Improved description for 3F_2 , 1F_3 & 3F_3

✓ Relativistic corrections sizeable !

Phase shifts-G wave





✓ Independent of renormalization scale

✓ Quantitatively better

✓ Relativistic effects small

Phase shifts-H wave & I wave



✓ High partial wave: **Two-pion exchange insignificant!**

Summary & outlook

□ Summary

- ✓ Calculate **two-pion exchange** contributions in **covariant BChPT**
- ✓ **Quantitatively better** description is achieved especially for **F wave**

□ Outlook

- ✓ The role of **nucleon resonance**
 - $\Delta(1232)$
 - **Roper resonance**

Thank you very much for your attention