HIDDeN
Hunting Invisibles: Dark sectors, Dark matter and Neutrinos

Neutrino upscattering to HNLs as an explanation of MB LEE

INVISIBLES '22

Jaime Hoefken Zink

In collaboration with A. Abdullahi, M. Hostert, D. Massaro, S. Pascoli

22/06/2022

This project has received funding / support from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska -Curie grant agreement No 860881-HIDDeN.

Anomaly: MiniBooNE Low Energy Excess



$560.6 \pm 119.6 (77.4 \pm 28.5)$ in $\nu(\bar{\nu})$ mode: $\mathbf{4.8\sigma}$

Anomaly: possible solutions

Topology	Model	Diagram	Signal
single γ	Transition magnetic moment		$N \to \nu\gamma$
	neutrino-induced inverse-Primakoff scattering		$a^* + A \to \gamma A$
e^+e^-	upscattering to N		$N \to \psi_{\text{inv}} e^+ e^-$
	bremsstrahlung of light Z'		$N^* \to \nu(Z' \to e^+e^-)$
	neutrino-induced Z' fusion		$Z'^* Z'^* \to (S \to e^+e^-)$
	neutrino-induced inverse-Primakoff scattering to Z'		$S^*\gamma^* \to (Z' \to e^+e^-)$
$\gamma\gamma$	neutrino-induced Z' fusion		$Z'^* Z'^* \to (S \to e^+e^-)$

Dark Photon Model

Portals



Dark photon model:

1. $U(1)_{\text{dark}}$ extension
2. dark Higgs singlet
3. ν_S and ν_D

Dark Photon Model

Misreconstruction at MiniBooNE



Condition: $< 13^\circ$ or $E_\nu < 0.03$ GeV





Simulation and Fitting

DarkNews



$$-2 \ln L =$$
$$\sum_{i,j} (D_i - P_i) M_{ij}^{-1} (D_j - P_j)$$
$$+ \ln \det M$$

Results

3+1



Best Fit: $m_{Z'}$=261.7MeV, m_4=223.3MeV,

χ^2_{red}=1.16, $|U^{\text{max}}_{\mu 4}|=10^{-2}$, $\epsilon=8\times10^{-4}$



Results

3+2



Best Fit: $m_{Z'}$=1.25GeV, m_5=107.5MeV, m_5=72.07MeV,

χ^2_{red}=0.91, $|U^{\text{max}}_{\mu 5(4)}|$=$10^{-3}$, ϵ=10^{-2}



Results

3+2 best fit



Summary

1. The 3+2 dark sector model can explain MB LEE in the off-shell decay regime of the HNL
2. It is a simple model with small number of parameters
3. The SBN program will be able to test it
4. It is not so constrained as other models, such as 3+1 DS

THANK YOU!

Extra Material

Lagrangian

$$\begin{aligned}\mathscr{L} =& \mathscr{L}_{\mathrm{SM}} + \overline{\nu}_D i \not{D}^x \nu_D \\ &+ (D^x_\mu \Phi)^\dagger (D^{x\mu} \Phi) - V(\Phi, H) \\ &- \frac{1}{4} X_{\mu\nu} X^{\mu\nu} - \frac{\sin\chi}{2} B_{\mu\nu} X^{\mu\nu} \\ &+ \overline{N} i \not{\partial} N - [y^\alpha_\nu (\overline{L}_\alpha \cdot \widetilde{H}) N^C + \frac{\mu'}{2} \overline{N} N^C + y_N \overline{N} \nu^C_D \Phi + \mathrm{h.c.}]\end{aligned}$$

$$\mathscr{L} \subset \epsilon e Z'_\mu J^\mu_{\mathrm{EM}} + g_D U_{Di} U^*_{Dj} Z'_\mu \left(i \overline{\nu}_i \not{\partial} \nu_j \right) \qquad (1)$$





Extra Material

MiniBooNE detector

- **Material**: pure mineral oil (CH_2) in spherical shape (radius: 6.1 m)
- **Baseline**: 541 m
- **Mean** energy: 800 MeV
- **Beam**: ν_μ, $\bar{\nu}_\mu$ (mainly), from BNB
- **Peaks**: 600 MeV (ν_μ), 400 MeV ($\bar{\nu}_\mu$)
- **Detection**: Cherenkov, PMTs



Extra Material

Neutrino Mixing

$$\mathscr{L}^{\nu}_{\text{mass}} = -\frac{1}{2}\begin{pmatrix}\overline{\nu_\alpha} & \overline{N} & \overline{\nu_D}\end{pmatrix}\begin{pmatrix}0_{3\times 3} & m_D^{\mathrm{T}} & 0\\ m_D & \mu' & \Lambda^{\mathrm{T}}\\ 0 & \Lambda & 0\end{pmatrix}\begin{pmatrix}\nu_\alpha\\ N^C\\ \nu_D^C\end{pmatrix} \tag{2}$$

$$\begin{pmatrix}\nu_e\\ \nu_\mu\\ \nu_\tau\\ N^C\\ \nu_D^C\end{pmatrix} \xrightarrow[\text{mass basis}]{\text{From flavor to}} \begin{pmatrix}\nu_1\\ \nu_2\\ \nu_3\\ \nu_4\\ \nu_5\end{pmatrix} \tag{3}$$

Extra Material

Couplings vs mass, 3+1



Extra Material

Couplings vs mass, 3+2



Extra Material

Signal efficiency