Higgs Hunting 2022



ID de Contribution: 80

Type: Non spécifié

Search for Lorentz boosted nonresonant HH production in the 4b final state with the CMS detector (Zoom)

mardi 13 septembre 2022 16:30 (12 minutes)

Double Higgs boson production (HH) allows us to measure the Higgs self-interaction and is uniquely sensitive to the structure of the Higgs potential. This talk will cover the production of HH \rightarrow 4b with highly boosted Higgs bosons in the gluon-fusion (ggF) and VBF production mode with 138 fb^{-1} of data collected with the CMS Experiment at sqrt(s) = 13 TeV [1]. The four-bottom-quark final state has the largest branching ratio (33.9%) amongst all HH decays, but is dominated by large backgrounds (QCD and top) and a poor decay channel resolution. To enhance the signal sensitivity, this analysis uses a dedicated jet identification algorithm developed to identify boosted H \rightarrow bb jets, known as the ParticleNet Tagger. This analysis is very sensitive to anomalous quartic VVHH couplings and, for the first time, we exclude $\kappa_{2V} = 0$ at >5 *sigma*, when other Higgs couplings are at their SM values. This talk will cover the analysis methods used and show the final sensitivity to different HH couplings (κ_{λ} , κ_{2V} and κ_V). I'll also discuss future directions and scope of improvements for measuring the HH process.

Orateur: DUTTA, Irene (California Institute of Technology (US)) **Classification de Session:** Young scientist forum