



ID de Contribution: 272

Type: Poster

Gate tunable spin-orbit coupling and superconductivity in KTaO₃-based two-dimensional electron gases

Oxide heterostructures are regarded as promising platforms for the next generation of electronic devices, displaying straightforward properties like high carrier mobility to exotic functionalities like multiferroicity. In polar heterostructures, the interfacial electric field can produce a two dimensional electron gas (2DEG). The broken inversion symmetry results in Rashba spin-orbit coupling that the source of rich physics. For example, it promotes the direct and inverse Edelstein effects that enable interconversion between spin and charge currents for spin-orbitronics. Furthermore, if the system is superconducting, a source of time-reversal symmetry breaking like a magnetic field could enable the generation of topological states.

The family of KTaO₃ 2DEGs are good candidates for such effects. On (111) and (110)-terminated KTaO₃, superconductivity is observed up to 2,2 K and 1 K respectively. The heavy Ta element confers a large Rashba coupling strength to the 2DEG, around 300 meV.Å for KTaO₃ (001), compared to few tens of meV.Å in LaAlO₃/SrTiO₃. However, to assess the functional perspectives of these systems, a fine control of their transport properties needs to be achieved.

Here, we present the tuning of superconductivity and spin-orbit coupling by field-effect in KTaO₃-based devices and relate to their band structure. The investigation of the interplay between these effects may shed light on the unconventional pairing mechanism.

Affiliation de l'auteur principal

UMPhy CNRS-Thales, LPEM ESPCI

Auteur principal: WITT, Hugo

Orateur: WITT, Hugo

Classification de Session: Session Poster 1: MC3, MC5, MC6, MC11, MC13, MC15, MC16, MC18, MC19, MC25, REDP, posters hors MC

Classification de thématique: MC19 Hétérostructures et interfaces de basse dimensionnalité