

21cm Intensity Mapping: opportunities and challenges on the road to the SKA Observatory

Marta Spinelli

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich



UNIVERSITY of the
WESTERN CAPE



Hydrogen through cosmic time

~380 kyr
Recombination
($z \sim 1100$)


200 Myr - 1 Gyr
Reionization
 $6 < z < 20$

~200 Myr
Cosmic Dawn
 $z \sim 20$


late Universe
Galaxy evolution
 $z < 6$

credit: ESA

Hydrogen through cosmic time



Hydrogen through cosmic time




21cm Cosmology


- signal *redshifted* due to the expansion of the Universe to **Radio Frequencies**
- **SKA Observatory**: cover **all the relevant frequencies** with unprecedented sensitivity



SKA-Low
50 MHz - 350 MHz
 $30 > z > 3$



SKA-Mid
350 MHz - 13.5 GHz
 $3 > z > 0$




credit: skatelescope.org

21cm Cosmology

- ❑ signal *redshifted* due to the expansion of the Universe to **Radio Frequencies**
- ❑ **SKA Observatory**: cover **all the relevant frequencies** with unprecedented sensitivity




SKA-Low
50 MHz - 350 MHz
 $30 > z > 3$




credit: skatelescope.org

SKA-Mid
350 MHz - 13.5 GHz
 $3 > z > 0$



MeerKAT
64 antennas
 $1.5 > z > 0$

Mapping neutral hydrogen



What is the nature of dark matter and dark energy?


- ❑ how is dark matter distributed on large scales?
- ❑ how does its distribution evolve with cosmic time?
- ❑ what is the role of dark energy?

How do baryons trace dark matter?

- ❑ what is the link between galaxies and dark matter halos?
- ❑ how are HI galaxies distributed in the cosmic web?
- ❑ how does the total cosmic HI evolve with redshift?

Intensity Mapping

credit: A. Pourtsidou




The distribution of **neutral Hydrogen** is a biased tracer of the **matter clustering** *similar to galaxy surveys*

In cosmology, **large scales** are fundamental

Intensity Mapping

credit: A. Pourtsidou




The distribution of **neutral Hydrogen** is a biased tracer of the **matter clustering**
similar to galaxy surveys

In cosmology, **large scales** are fundamental

How can we efficiently observe cosmological volumes?

Intensity Mapping

credit: A. Pourtsidou



The distribution of **neutral Hydrogen** is a biased tracer of the **matter clustering** *similar to galaxy surveys*


In cosmology, **large scales** are fundamental

How can we efficiently observe cosmological volumes?

Intensity Mapping: total intensity of the 21 cm emission line in a **large pixel** (low spatial resolution)

Intensity Mapping

credit: A. Pourtsidou




The distribution of **neutral Hydrogen** is a biased tracer of the **matter clustering** *similar to galaxy surveys*

Intensity Mapping: total intensity of the 21cm emission line in a **large pixel** (low spatial resolution)

different frequencies,
different z
high spectral resolution
(tomography)

Key cosmological probe




Key cosmological probe




HIP-POP
MS et al. 2022

Key cosmological probe

SKA Red Book (2020)



SKA Red Book (2020)



Berti, MS et al. 2022

Maria's talk yesterday!

SKAO forecasts

$$P_{21}(z, k, \mu) = \bar{T}_b^2(z) \left[b_{\text{HI}}(z) + f(z) \mu^2 \right]^2 P_m(z, k)$$

SKAO forecasts

$$P_{21}(z, k, \mu) = \bar{T}_b^2(z) \left[b_{\text{HI}}(z) + f(z) \mu^2 \right]^2 P_m(z, k)$$

$$P_\ell(z, k) = \frac{(2\ell + 1)}{2} \bar{T}_b^2(z) P_m(z, k) \int_{-1}^1 d\mu \mathcal{L}_\ell(\mu) \left[b_{\text{HI}}(z) + f(z) \mu^2 \right]^2$$


SKAO forecasts

$$P_{21}(z, k, \mu) = \bar{T}_b^2(z) \left[b_{\text{HI}}(z) + f(z) \mu^2 \right]^2 P_m(z, k)$$

$$P_\ell(z, k) = \frac{(2\ell + 1)}{2} \bar{T}_b^2(z) P_m(z, k) \int_{-1}^1 d\mu \mathcal{L}_\ell(\mu) \left[b_{\text{HI}}(z) + f(z) \mu^2 \right]^2$$


- ❑ We consider only monopole and quadrupole $l=0,2$
- ❑ SKA-Mid like observations
 - **tomographic** (6 redshift between 0 and 3)
 - **Single-dish**: beam effect
 - expected noise and sky area

Berti, MS et al. 2022
[arXiv:2209.07595](https://arxiv.org/abs/2209.07595)





SKAO forecasts

Berti, MS et al. 2022




SKAO forecasts

Berti, MS et al. 2022




The challenge of foregrounds




adapted from Haslam et al. (1982)

The challenge of foregrounds




adapted from Haslam et al. (1982)

The challenge of foregrounds




adapted from Haslam et al. (1982)

The challenge of foregrounds




adapted from Haslam et al. (1982)

The challenge of foregrounds




The challenge of foregrounds





Properties of the foregrounds

Matshawule, MS et al. 2021





Properties of the foregrounds

Matshawule, MS et al. 2021



Properties of the foregrounds



Matshawule, MS et al. 2021



- ❑ foregrounds are orders of magnitude **stronger** than the 21cm signal
- ❑ their frequency behaviour is **smooth** (highly correlated)

Properties of the foregrounds

Matshawule, MS et al. 2021




- ❑ foregrounds are orders of magnitude **stronger** than the 21cm signal
- ❑ their frequency behaviour is **smooth** (highly correlated)

Questions:


- ❑ Can the **properties of the foregrounds** be used to separate them from the pristine **21cm signal**?
- ❑ **Even if we add some realism to our simulations?** (beam response, noise, RFI, polarization leakage, ...)

Dealing with foregrounds

Various strategies: e.g. **modelling**, **avoidance** and **separation/cleaning**



Bowman et al. (2018)




Liu et al. (2014)

separation/cleaning:
PCA, kPCA, FastICA,
GMCA, GPR, ML-GPR

see also Isabella's talk yesterday!

A cleaning example

Mock observation “cube”



Simulation includes:

- ❑ 100 channels around redshift 0.5
- ❑ Foreground contamination:
Synchrotron, Free-free, point sources
- ❑ Gaussian beam
- ❑ White noise

$$T = As + n + c$$

A mixing matrix of the foreground sources

noise


Cosmological signal

How many sources?


N_{fg} need to be estimated/guessed

A cleaning example



S map = 0










S map = 1



S map = 2



A cleaning example





A cleaning example

$$\mathbf{c} + \mathbf{n} = \mathbf{T} - \mathbf{A}\mathbf{s}$$




A mixing matrix including only the first Nfg components





With GBT data

Wolz et al. 2022






With GBT data

Wolz et al. 2022



Mitigation of systematics with cross-correlation



Auto Correlation:

$$\langle \mathbf{X}_{\text{opt}} \mathbf{X}_{\text{opt}} \rangle = \langle \mathbf{S}_{\text{opt}} \mathbf{S}_{\text{opt}} \rangle + 2 \langle \mathbf{S}_{\text{opt}} \mathbf{N}_{\text{opt}} \rangle + \langle \mathbf{N}_{\text{opt}} \mathbf{N}_{\text{opt}} \rangle$$

uncorrelated

$$\langle \mathbf{X}_{\text{opt}} \mathbf{X}_{\text{opt}} \rangle = \langle \mathbf{S}_{\text{opt}} \mathbf{S}_{\text{opt}} \rangle + \langle \mathbf{N}_{\text{opt}} \mathbf{N}_{\text{opt}} \rangle$$

signal you want

noise you don't want

Cross Correlation:



$$\langle \mathbf{X}_{\text{opt}} \mathbf{X}_{\text{rad}} \rangle = \langle \mathbf{S}_{\text{opt}} \mathbf{S}_{\text{rad}} \rangle + \langle \mathbf{S}_{\text{opt}} \mathbf{N}_{\text{rad}} \rangle + \langle \mathbf{S}_{\text{rad}} \mathbf{N}_{\text{opt}} \rangle + \langle \mathbf{N}_{\text{opt}} \mathbf{N}_{\text{rad}} \rangle$$

courtesy of Steve Cunnington

Intensity Mapping with MeerKAT




Antennas	All 64 MeerKAT dishes
Observation mode	Single-dish
Frequency range	0.856-1.712 GHz
Frequency resolution	0.2 MHz
Time resolution	2s
Exposure time	1.5hr x 7 scans
Target field	WiggleZ 11hr field ($10^\circ \times 30^\circ$)



MeerKAT observations


Wang et al. 2021
[including MS]



MeerKLASS: 64 MeerKAT antennas used in **single-dish mode**



- ❑ first successful calibration of **intensity mapping data from MeerKAT**
- ❑ L-band: 850-1700 MHz (4096 channels)

MeerKLASS results





Cunnington et al. 2022 [including MS]

MeerKLASS results



Cunnington et al. 2022 [including MS]

MeerKLASS results



Cunnington et al. 2022 [including MS]

Towards the SKA Observatory

We have:

21cm intensity mapping data difficult to clean (signal only in cross-correlation)

Simulations that are still not a realistic representation of the actual data

Cleaning methods that have still to be extensively tested with realistic simulations

We would like:

More and better data

More realistic simulations mimicking the data


More sophisticated cleaning methods tested on more realistic simulations

Final aim:

A 21cm **(auto) power spectrum detection** validated with realistic simulations and tested with various and robust cleaning methods

Ingredients for the simulations



Credit: D. Alonso



Mock 21cm maps

HI properties




- HI is a key ingredient for galaxy evolution
- estimation of HI bias properties and of the **$M_{\text{HI}}-M_{\text{halo}}$ relation** using the semi-analytical model **GAEA** (the **G**ALaxy **E**volution and **A**ssembly model, De Lucia et al. 2016, Xie et al. 2018)



Mock 21cm maps

Fast 21cm intensity map generation

- fundamental for **cosmological volume** and **end-to-end simulations**
- HOD methods on fast halo catalogues
MS et al. 2022



Foregrounds


Typical modeling:

Haslam 408 MHz
Ramazeilles et al. (2015)




Spatially varying synch spectral index
Miville-Deschenes et al. (2008)

Free-Free from Planck Sky Model
Delabruille et al (2013)

Extragalactic PS
Olivari (2018), Matshwule et al. (2021)



Instrumental effects



Need a realistic beam modeling
side-lobes, frequency evolution,
more accurate deconvolution


Scanning strategy
non homogeneous noise, need
for real space convolution,
polarization leakage

Radio Frequency Interference (RFI)
impact on cleaning,
impact on signal interpretation


Instrumental effects: the beam

Airy beam
Harper et al.
(2018)

- ❑ MeerKAT beam has **side-lobes** (same for SKA-MID)
- ❑ a strong point source in the side-lobes contaminates the signal and **can complicate the foreground subtraction**




Eidos: measured
Asad et al. 2020



The beam evolves with frequency

Effect of the telescope beam



Matshawule, MS et al. 2021

a realistic **MeerKAT** beam: side-lobes (cosine) and a non-trivial frequency evolution (ripple)

- ❑ **point sources** and synchrotron spatial structures coupled with the beam **complicate the cleaning**
- ❑ Careful **beam-deconvolution** alleviates the problem but need to be careful for precision cosmology

Foreground subtraction challenge


(subset of) SKA IM Focus Group

Project setup:

- ❑ various foreground models and realistic HI maps
- ❑ **instrumental modeling MeerKAT-like and SKAO-like**
- ❑ 9 different foreground removal methods (PCA, FastICA, ...)

Blind challenge to discover weaknesses and strengths of the various methods

Isabella Carucci, Steve Cunnington, Ze Fonseca, Stuart Harper, Mel Irfan, Alkistis Pourtsidou, Marta Spinelli, Laura Wolz





given IM data now, would your favorite method extract the cosmological signal?

Foreground subtraction challenge

- ❑ How much can **instrument/foregrounds coupling** impact the signal reconstruction?
- ❑ definition of statistics and metrics to evaluate the relative performances

Realistic instrumental effects inevitably complicate the foreground cleaning



MS et al. 2022

Moving forward

Data:



Keep working with pathfinder data (MeerKLASS) to understand the instrument and improve the pipelines

Simulations:


Improve and refine end-to-end simulations

Final aim:

A 21 cm **(auto) power spectrum detection** validated with realistic simulations



- ❑ new L band data under analysis (41 x 1.5 h)
- ❑ UHF band available (could go to higher redshift)



Conclusions

- ❑ **21cm Cosmology** still have to prove its full potential but offers an incredible window into the evolution of the Universe
- ❑ Intensity Mapping surveys are taking data (and new instrument are planned)
- ❑ **detection in cross-correlation** from MeerKLASS survey x galaxy survey (7.7σ)
- ❑ analysing new data: effort in understanding the instrument and developing better analysis pipelines
- ❑ Keep improving the simulations: both signal, foregrounds and instrumental effects
- ❑ **Prepare for the SKAO era and its potential contribution to the knowledge of large-scale structures**