

Search for extra Higgs bosons in same sign top-quark pair + jets final state

Ting-Hsiang Hsu (National Taiwan University)

On behalf of the CMS collaboration


**Higgs Hunting Conference 2023
Sep 11 2023**



Introduction to Extra Higgs Bosons

- Standard model still has several unsolved puzzles i.e. (anti)matter imbalance, $g-2$ muon anomaly etc.
- Several models introduce an extended Higgs sector i.e. supersymmetry (SUSY), Two-Higgs-Doublet Model (2HDM) in different variations etc.
- The second Higgs doublet in a 2HDM-like model introduces 5 scalar bosons
 - Charged scalar: H^\pm
 - Neutral scalar: H, h
 - Neutral pseudoscalar: A
- Many searches have been performed for extra Higgs Bosons

CMS Higgs Group Summary



FCNC in 2HDM

- **Glashow-Weinberg condition** is usually imposed in 2HDM to avoid flavor changing neutral currents (FCNC).
- At the **alignment limit**, in a 2HDM without Glashow-Weinberg condition (type-III 2HDM / g2HDM), FCNC of 125 GeV Higgs boson are suppressed but not for exotic Higgs.
 - **Exotic Higgs in subTeV** level and **O(1) extra Yukawa couplings** are not yet excluded.
 - **Muon g-2 anomaly** and **baryogenesis** can be explained by such models.


$$y_{ij} = \sqrt{2}m_i\delta_{ij}/v$$

$$q_i q_j h \text{ coupling (125 GeV Higgs) } \propto -y_{ij} \sin_\gamma + \rho_{ij} \cos_\gamma$$

$$q_i q_j H \text{ coupling (exotic Higgs) } \propto y_{ij} \cos_\gamma + \rho_{ij} \sin_\gamma$$

Alignment limit : $\cos_\gamma \rightarrow 0$


FCNC of exotic Higgs sector provides chance to observe/
exclude **subTeV** exotic Higgs and **O(1) extra Yukawa couplings**



Target Process

Target FCNC process

CMS-PAS-TOP-22-010



We test different mass $m_{H/A}$ and coupling strength ρ_{tu} & ρ_{tc} assumptions


Extra Yukawa Coupling

- Probe extra Yukawa coupling involving top quark (ansatz like $\rho_{qq'} \propto \sqrt{m_q m_{q'}}$)
- Probe ρ_{tu} & ρ_{tc} through $tt\bar{u}$ & $tt\bar{c}$ processes separately
- Probe range: $0.1 \leq \rho_{tq} \leq 1.0$ (assume all other extra Yukawa couplings to be 0)

Target Process

Target FCNC process

CMS-PAS-TOP-22-010



We test different mass $m_{H/A}$ and coupling strength ρ_{tu} & ρ_{tc} assumptions

Exotic Higgs Bosons

Two scenarios are considered

H and A non-interference


- Only consider A or H
- Probe range: $200 \text{ GeV} \leq m_{A/H} \leq 1 \text{ TeV}$

H and A interference

- Take $m_A - m_H = 50 \text{ GeV}$ for demo
- Probe range: $250 \text{ GeV} \leq m_A \leq 1 \text{ TeV}$

Object & Event Selection

Target FCNC process



$$ug/cg \rightarrow tH \rightarrow tt\bar{u}/t\bar{c} \rightarrow (l^+b\nu)(l^+b\nu)\bar{u}/\bar{c}$$

Signal Topology

Two same sign leptons
At least three jets
Missing transverse momentum

Signal cross section

Depending on the coupling value, mass and the H-A interference assumption


ttu processes range from $\sim 1 \times 10^{-4}$ to $\sim 7 \times 10^{-1} pb$

ttc processes range from $\sim 5 \times 10^{-6}$ to $\sim 7 \times 10^{-2} pb$

CMS-PAS-TOP-22-010

Background Contribution



(Before BDT cut)



Analysis Technique

- DeepJet Flavor Tagging

- Neural network utilizes global variables, charged/neutral particle and secondary vertex features in the jets to perform flavor tagging.




Analysis Technique

- Boosted Decision Tree

- Input: Jet flavor info, lepton/jet kinematic, MET, HT, and object pair kinematic information
- BDT > -0.6 cut is imposed in order to improve the stability of the fit and the corresponding fit uncertainties

Input features of the BDT

$p_T(\ell_i): i=1,2; H_T, p_T^{\text{miss}}$
 $CvsB(j_i), CvsL(j_i): i=1,2,3$
 $m_{\ell\ell}, m_{\ell\ell}(j_i): i=1,2,3$
 $\Delta R(j_n, j_m), m(j_n, j_m): 1 \leq n < m \leq 3$
 $\Delta R(j_n, l_m), m(j_n, l_m): n=1,2,3; m=1,2$





Uncertainties

- Fit is performed using BDT distribution.
- Profiled uncertainty is used for analysis.
- Main uncertainties come from
 - Statistical unc. from the fit
 - Flavor tagging
 - Nonprompt background estimation
 - ttW cross section

Post-fit uncertainty sources with respect to the total uncertainty
(given in percentages)

	$\rho_{tu} = 0.4$	
	$m_A = 200 \text{ GeV (1 TeV)}$ Without interference	$m_A = 250 \text{ GeV (1 TeV)}$ With interference
Jet flavor identification	16 (55)	26 (46)
Nonprompt leptons	39 (13)	43 (13)
ttW background	33 (28)	35 (30)
Total signal modeling	20 (16)	7 (19)
Total background modeling	67 (56)	65 (59)
Total experimental	44 (56)	55 (50)
Statistical	57 (57)	52 (61)
	$\rho_{tc} = 0.4$	
Jet flavor identification	18 (47)	24 (48)
Nonprompt leptons	35 (14)	31 (14)
ttW background	32 (32)	34 (31)
Total signal modeling	14 (12)	16 (13)
Total background modeling	66 (62)	64 (61)
Total experimental	44 (50)	46 (52)
Statistical	61 (61)	60 (59)




Results

- Probe phase space with O(1) extra Yukawa couplings and sub TeV exotic Higgs mass
- Results are consistent with standard model predictions.

Observed (expected) lower limits on $m_A(m_H)$ at 95% CL.

Observed (expected) mass limit [GeV]			
ρ_{tu}	Without interference m_A or m_H	With interference m_A	With Interference m_H
0.1	— (—)	— (—)	— (—)
0.4	900 (900)	1000 (1000)	950 (950)
1.0	1000 (1000)	1000 (1000)	950 (950)
ρ_{tc}	m_A or m_H	m_A	m_H
0.1	— (—)	— (—)	— (—)
0.4	— (—)	300 (350)	250 (300)
1.0	700 (600)	800 (550)	650 (500)

Observed and expected 95% CL upper limit on the signal strength




Summary


- Search for top quark related flavor changing neutral current in extended Higgs sector using full run2 CMS data
- Good agreement with standard model predictions.
- Exclude certain phase space of O(1) extra Yukawa couplings and sub TeV exotic Higgs mass with/without considering H/A interference.

CMS-PAS-TOP-22-010

Thank you for your attention



95% CL upper limit on $\mu = \sigma(\text{obs})/\sigma(\text{theory})$



95% CL upper limit on $\mu = \sigma(\text{obs})/\sigma(\text{theory})$

Back up

Previous CMS results

- Many researches have been performed regarding to extra Higgs Bosons

$H/A \rightarrow tt$: Eur. Phys. J. C 77 (2017) 578

$H/A \rightarrow bb$: JHEP 08 (2018) 113

$H/A \rightarrow \tau\tau$: JHEP 09 (2018) 007

$H/A \rightarrow \mu\mu$: Phys. Lett. B 798 (2019)

$A \rightarrow Zh \rightarrow (ll, \nu\nu)bb$: Eur. Phys. J. C 79 (2019)

$H/A \rightarrow Z(ll)A/H(bb)$: JHEP 03 (2020) 055

$H \rightarrow WW$: JHEP 03 (2020) 034

$X \rightarrow YH \rightarrow b\bar{b}b\bar{b}$: Phys. Lett. B 842 (2023) 137392

$\phi \rightarrow \tau\tau$: JHEP 07 (2023) 073

$H \rightarrow AA \rightarrow 4\gamma$: Accepted pub. In Phys. Rev. Lett.

$H \rightarrow e\mu$: Submitted to Phys. Rev. D

$H \rightarrow \gamma\gamma$: CMS-PAS-HIG-20-002

$\phi \rightarrow ll$: CMS-PAS-EXO-21-018

$H^\pm \rightarrow \tau_h\nu$: JHEP 2019:142

$H^\pm \rightarrow Wa$: Phys. Rev. Lett. 123 131802 (2019)

$H^\pm \rightarrow tb$: JHEP 2020:096, JHEP 2020:126

$H^\pm \rightarrow cs, cb$: Phys. Rev. D 102, 072001 (2020)



$H^\pm \rightarrow H(\tau\tau)W$: Accepted pub. In JHEP

FCNC in extended Higgs sector still remains to be studied.

Analysis Technique - DeepJet

arXiv:2008.10519

- DeepJet Flavor Tagging
 - Neural network utilizing global variables, charged/neutral particle and secondary vertex features in the Jets to perform flavor tagging.




BDT distribution


- Boosted Decision Tree

- Input: Jet flavor info, lepton/jet kinematic, MET, HT, and object pair kinematic information
- BDT > -0.6 cut is imposed in order to improve the stability of the fit and the corresponding fit uncertainties

BDT focuses on ttu



BDT focuses on ttc



Results

CMS-PAS-TOP-22-010

H-A non-interference
Only consider ρ_{tu}

H-A non-interference
Only consider ρ_{tc}

H-A interfere
Only consider ρ_{tu}





H-A interfere
Only consider ρ_{tc}

CMS Preliminary 138 fb⁻¹ (13 TeV)

CMS Preliminary 138 fb⁻¹ (13 TeV)

CMS Preliminary 138 fb⁻¹ (13 TeV)

CMS Preliminary 138 fb⁻¹ (13 TeV)



Results - 2D phase space

CMS-PAS-TOP-22-010


H-A non-interference
Only consider ρ_{tu}

H-A non-interference
Only consider ρ_{tc}

H-A interfere
Only consider ρ_{tu}


H-A interfere
Only consider ρ_{tc}

CMS Preliminary 138 fb⁻¹ (13 TeV)




95% CL upper limit on $\mu=\alpha(\text{obs})/\alpha(\text{theory})$

CMS Preliminary 138 fb⁻¹ (13 TeV)




95% CL upper limit on $\mu=\alpha(\text{obs})/\alpha(\text{theory})$

CMS Preliminary 138 fb⁻¹ (13 TeV)



95% CL upper limit on $\mu=\alpha(\text{obs})/\alpha(\text{theory})$

CMS Preliminary 138 fb⁻¹ (13 TeV)



95% CL upper limit on $\mu=\alpha(\text{obs})/\alpha(\text{theory})$

Background Category

Category	Samples	Category	Samples
TT	TTTo2L	Others	tW & tbarW DY ttZZ ttWW ttWZ ttWH ttZH ttZ(ll + qq) tZq tttj tttW ttt ZZZ WZZ WWZ WWW
VV	WW(OS) WZ(QCD)		
VBS	WpWpJJ(EWK+QCD) WLLjj ZZJJTo4L		
ttH	ttH		
ttW	ttWtoLnu ttWtoQQ		