

SÉMINAIRE du PÔLE THÉORIE



LABORATOIRE DE PHYSIQUE DES DEUX
INFINIS IRÈNE JOLIOT-CURIE
Pôle de Physique Théorique
Bât. 100, F-91406 ORSAY CEDEX
Tél (33)-(0)1-6915-7330 - Fax (33)-(0)1-6915-7748

université
PARIS-SACLAY

Tobias Frederico

Instituto Tecnológico de Aeronautica

Two-neutrons halos and beyond

Weakly bound halos formed in unstable exotic nuclei at the drip line is by now a well grounded description. In the case of a core plus two-neutrons, for example ${}^6\text{He}$, ${}^{11}\text{Li}$, ${}^{14}\text{Be}$, ${}^{19}\text{B}$, ${}^{22}\text{C}$, which due to the dilute halo allows its understanding in terms of universal properties, quite independent of the nuclear model and dominated by few physical scales, like the large scattering lengths/low-lying resonances and the two-neutron separation energy. The next challenge is neutron halos with more neutrons, which will be qualitatively illustrated by representative cases, like e.g. ${}^{28}\text{O}$, where again few physical scales should be dominant on the properties of nuclei like that.

Thursday 12 October 2023,

14h00

IJCLab, Build. 100, Room A018